

Durham Research Online

Deposited in DRO:

22 May 2020

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Budge, Lucy and Campbell, John M. and De Laurentis, Giuseppe and Ellis, R. Keith and Seth, Satyajit (2020) 'The one-loop amplitudes for Higgs + 4 partons with full mass effects.', *Journal of high energy physics.*, 2020 (5). 079.

Further information on publisher's website:

[https://doi.org/10.1007/JHEP05\(2020\)079](https://doi.org/10.1007/JHEP05(2020)079)

Publisher's copyright statement:

This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

The one-loop amplitudes for Higgs + 4 partons with full mass effects

Lucy Budge,^a John M. Campbell,^b Giuseppe De Laurentis,^a R. Keith Ellis^a
and Satyajit Seth^c

^a*Institute for Particle Physics Phenomenology, Durham University,
Durham, DH1 3LE, U.K.*

^b*Fermilab,
PO Box 500, Batavia IL 60510-5011, U.S.A.*

^c*Physical Research Laboratory,
Navrangpura, Ahmedabad — 380009, India*

E-mail: lucy.budge@durham.ac.uk, johnmc@fnal.gov,
giuseppe.de-laurentis@durham.ac.uk, keith.ellis@durham.ac.uk,
seth@prl.res.in

ABSTRACT: We present compact analytic formulae for the one-loop amplitudes for Higgs + 4 parton scattering, $0 \rightarrow ggggh$, $0 \rightarrow \bar{q}qggh$ and $0 \rightarrow \bar{q}q\bar{q}'q'h$, mediated by a loop of massive coloured quarks. We exploit the correspondence with a theory in which a massive coloured scalar circulates in the loop to avoid a proliferation in the number of terms in the result. In addition, we use momentum twistors and high precision numerical evaluations to simplify the expressions. The analytic results in this paper, in terms of spinor products, allow construction of an efficient numerical program to calculate the amplitude.

KEYWORDS: NLO Computations

ARXIV EPRINT: [2002.04018](https://arxiv.org/abs/2002.04018)

Contents

1	Introduction	1
2	Structure of the calculation	2
2.1	Definition of colour amplitudes	2
2.2	Decomposition into scalar integrals	4
2.3	Unitarity methods	6
2.4	Simplification techniques	6
3	Higgs boson production mediated by a coloured scalar	8
3.1	Relationship of the fermion theory to the scalar theory	9
4	Coefficients for $H_4^{1234}(g^+, g^+, g^+, g^+; h)$	11
4.1	Boxes	12
4.2	Triangles	13
4.3	Rational terms	13
5	Coefficients for $H_4^{1234}(g^+, g^+, g^+, g^-; h)$	13
5.1	Boxes	15
5.2	Triangles	17
5.3	Bubbles	18
5.4	Rational terms	19
6	Coefficients for $H_4^{1234}(g^+, g^-, g^+, g^-; h)$	19
6.1	Boxes	20
6.2	Triangles	21
6.3	Bubbles	23
6.4	Rational terms	23
7	Coefficients for $H_4^{1234}(g^+, g^+, g^-, g^-; h)$	24
7.1	Boxes	24
7.2	Triangles	25
7.3	Bubbles	27
7.4	Rational terms	27
8	Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^+, g^+; h)$	28
8.1	Boxes	28
8.2	Triangles	29
8.3	Bubbles	30
8.4	Rational terms	30

9	Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^-, g^+; h)$	30
9.1	Boxes	30
9.2	Triangles	31
9.3	Bubbles	32
9.4	Rational terms	33
10	Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^+, g^-; h)$	33
10.1	Triangle	33
10.2	Bubble	34
10.3	Rational terms	34
11	Amplitude for $0 \rightarrow \bar{q}q\bar{q}qh$	34
12	Large mass limit	35
13	Conclusions	36
A	Integrals	37
B	Spinor algebra	37
C	Results for tree-level amplitudes with massive scalars	38
C.1	One gluon	39
C.2	Two gluons	39
C.3	Three gluons	39
C.4	One gluon, two quarks	40
D	Numerical value of coefficients at a given phase-space point	40

1 Introduction

At the Large Hadron Collider (LHC) the primary mechanism for producing and detecting Higgs bosons is the process $gg \rightarrow h$. This process is mediated, in the Standard Model, by a loop of massive coloured fermions. Since the Yukawa coupling is proportional to the fermion mass the predominant contribution is the result of the coupling of the top quark to the Higgs boson. In the limit in which only a very heavy top quark contributes, the corresponding amplitude is independent of the top quark mass; this gives rise to an effective field theory (EFT) in which the loop of heavy top quarks is replaced by an effective Lagrangian,

$$\mathcal{L}_{\text{eff}} = \frac{g_s^2}{48\pi^2 v} h G_{\mu\nu}^A G^{A,\mu\nu} \quad (1.1)$$

where g_s is the strong coupling constant, v is the vacuum expectation value of the Higgs field, $G_{\mu\nu}$ is QCD field strength, and h is the Higgs boson field. The EFT in eq. (1.1)

has been used to compute higher-order corrections to the inclusive cross-section — most recently up to next-to-next-to-next-to-leading order [1, 2] — as well as rates for the production of Higgs bosons in association with up to three additional jets up to next-to-leading order [3, 4]. The effective field theory description is expected to break down when, for example, the transverse momentum of produced gluons is of order of the top quark mass. This breakdown has most recently been investigated at NLO in ref. [5]. This kinematic regime is beginning to be explored at the LHC [6] and can give important information about the mediators in the loop that couple to the Higgs. For such configurations it is therefore important to make use of a superior calculation in which the full dependence on the top quark mass is retained. Such a calculation also allows a direct quantification of the breakdown of the EFT approach.

Analytic results for the Higgs+3 parton amplitude in the full theory have been known for a long time [7, 8]. Corresponding results for Higgs+4 parton amplitudes have been obtained in refs. [9, 10], although in both cases expressions for at least some of the amplitudes were too long to report. In addition there are several automatic procedures than can provide numerical results for one-loop amplitudes [11–14]. The aim of this paper is to present compact amplitudes for all contributing processes,

$$0 \rightarrow ggggh, \quad (1.2)$$

$$0 \rightarrow \bar{q}qggh, \quad (1.3)$$

$$0 \rightarrow \bar{q}q\bar{q}'q'h, \quad (1.4)$$

retaining all mass effects. Compact analytic results for the $0 \rightarrow ggggh$ case when all the gluons have positive helicity have been published in ref. [15]. Although our result is therefore not new *per se*, it is the first time that a compact publishable analytic result has been obtained for all gluon helicities. A calculation with compact analytic formulae allows examination of the structure of the amplitude for all values of the fermion mass. It also has the potential to lead to faster and more stable numerical evaluation of the amplitude. This would be a boon to calculations requiring this amplitude in all regions of phase space, such as recent NLO predictions for Higgs boson plus 1-jet production in the full theory [5] and at large transverse momentum [16, 17]. Although the results are quite compact, given the number of integral coefficients, this paper is not easy to read. However we believe that it is detailed enough that readers wishing to implement this amplitude in a numerical program will find enough information to do so in our paper.

2 Structure of the calculation

2.1 Definition of colour amplitudes

The amplitude for the production of a Higgs boson and n gluons can be expressed in colour-ordered sub-amplitudes as follows:

$$\mathcal{H}_n^{gggg}(\{p_i, h_i, c_i\}) = i \frac{g_s^n}{16\pi^2} \frac{m^2}{v} \sum_{\{1,2,\dots,n\}'} \text{tr}(t^{c_1} t^{c_2} \dots t^{c_n}) H_n^{\{c_i\}}(1^{h_1}, 2^{h_2}, \dots, n^{h_n}; h), \quad (2.1)$$

where the sum with the *prime*, $\sum_{\{1,2,\dots,n\}'}$, is over all $(n-1)!$ *non-cyclic* permutations of $1, 2, \dots, n$ and the t matrices are the $SU(3)$ matrices in the fundamental representation normalized such that,

$$\text{tr}(t^a t^b) = \delta^{ab}. \quad (2.2)$$

m is the mass of the quark circulating in the loop. Because of Bose symmetry it is sufficient to calculate one permutation, and the other colour sub-amplitudes can be obtained by exchange.

For the particular case at hand with four gluons eq. (2.1) becomes,

$$\begin{aligned} \mathcal{H}_4^{gggg}(\{p_i, h_i, c_i\}) = & i \frac{g_s^4}{16\pi^2} \left(\frac{m^2}{v} \right) \left\{ [\text{tr}(t^{c_1} t^{c_2} t^{c_3} t^{c_4}) + \text{tr}(t^{c_1} t^{c_4} t^{c_3} t^{c_2})] H_4^{1234}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \right. \\ & + [\text{tr}(t^{c_1} t^{c_3} t^{c_4} t^{c_2}) + \text{tr}(t^{c_1} t^{c_2} t^{c_4} t^{c_3})] H_4^{1342}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \\ & \left. + [\text{tr}(t^{c_1} t^{c_4} t^{c_2} t^{c_3}) + \text{tr}(t^{c_1} t^{c_3} t^{c_2} t^{c_4})] H_4^{1423}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \right\}. \end{aligned} \quad (2.3)$$

Squaring the amplitude eq. (2.3) for a fixed helicity configuration and summing over colours we find

$$\begin{aligned} \sum_{\text{colours}} |\mathcal{H}_4^{gggg}|^2 = & \left[\frac{g_s^4}{16\pi^2} \left(\frac{m^2}{v} \right) \right]^2 (N^2 - 1) \left\{ 2N^2 (|H_4^{1234}|^2 + |H_4^{1342}|^2 + |H_4^{1423}|^2) \right. \\ & \left. - 4 \frac{(N^2 - 3)}{N^2} |H_4^{1234} + H_4^{1342} + H_4^{1423}|^2 \right\}, \end{aligned} \quad (2.4)$$

where N is the dimensionality of the $SU(N)$ colour group, i.e. $N = 3$, and the labels for the helicity configuration (as explicitly shown in eq. (2.3)) have been suppressed.

The amplitude for the production of a Higgs boson, an antiquark, quark and two gluons is similarly decomposed into colour-ordered amplitudes as follows,

$$\begin{aligned} \mathcal{H}_4^{\bar{q}ggg}(\{p_i, h_i, c_i, j_i\}) = & i \frac{g_s^4}{16\pi^2} \left(\frac{m^2}{v} \right) \left[(t^{c_3} t^{c_4})_{j_2 j_1} H_4^{34}(1^{h_1}, 2^{-h_1}, 3^{h_3}, 4^{h_4}; h) \right. \\ & \left. + (t^{c_4} t^{c_3})_{j_2 j_1} H_4^{43}(1^{h_1}, 2^{-h_1}, 3^{h_3}, 4^{h_4}; h) \right]. \end{aligned} \quad (2.5)$$

The colour structure $\delta^{c_3 c_4} \delta_{j_2 j_1} / N$ is also present in individual diagrams but makes no net contribution to the one-loop amplitude. In this paper we will give results for the colour-ordered amplitude H_4^{34} . It is straightforward to obtain H_4^{43} from this through the parity operation (complex conjugation) and permutation of momentum labels. Squaring and summing over colours yields,

$$\sum |\mathcal{H}_4^{\bar{q}ggg}|^2 = \left(\frac{g_s^4}{16\pi^2} \right)^2 \left(\frac{m^2}{v} \right)^2 (N^2 - 1) \left[N (|H_4^{34}|^2 + |H_4^{43}|^2) - \frac{1}{N} |H_4^{34} + H_4^{43}|^2 \right], \quad (2.6)$$

where the labelling of the helicity configuration shown in eq. (2.5) has again been suppressed.

The four-quark amplitude takes the form,

$$\mathcal{H}_4^{4q}(\{p_i, h_i, j_i\}) = i \frac{g_s^4}{16\pi^2} \left(\frac{m^2}{v} \right) (t^{c_1})_{j_2 j_1} (t^{c_1})_{j_4 j_3} H_4^{4q} \left(1_{\bar{q}}^{h_1}, 2_q^{-h_1}, 3_{\bar{q}'}^{h_3}, 4_{q'}^{-h_3} \right) \quad (2.7)$$

where the helicities of the quarks are fixed by those of the antiquarks. Performing the sum over colours we then have,

$$\sum |\mathcal{H}_4^{4q}(h_1, h_3)|^2 = \left(\frac{g_s^4}{16\pi^2}\right)^2 \left(\frac{m^2}{v}\right)^2 (N^2 - 1) |H_4^{4q}(h_1, h_3)|^2 \quad (2.8)$$

for the case in which the quark lines have different flavours. For the case of identical quarks we first introduce,

$$H_4^{4q'}(h_1, h_3) = H_4^{4q}(1_{\bar{q}}^{h_1}, 4_q^{-h_1}, 3_{\bar{q}}^{h_3}, 2_q^{-h_3}) \quad (2.9)$$

The sum over the colours for the identical case is then,

$$\begin{aligned} \sum |\mathcal{H}_4^{4q}|^2 &= \left(\frac{g_s^4}{16\pi^2}\right)^2 \left(\frac{m^2}{v}\right)^2 (N^2 - 1) \left(|H_4^{4q}(h_1, h_3)|^2 + |H_4^{4q'}(h_1, h_3)|^2 \right. \\ &\quad \left. + \frac{\delta_{h_1 h_3}}{N} \left(H_4^{4q}(h_1, h_3) H_4^{4q'}(h_1, h_3)^* + H_4^{4q}(h_1, h_3)^* H_4^{4q'}(h_1, h_3) \right) \right) \end{aligned} \quad (2.10)$$

where, as indicated, the term on the second line only contributes when the quarks have the same helicity.

2.2 Decomposition into scalar integrals

The colour-ordered sub-amplitudes can be expressed in terms of scalar integrals. For the $0 \rightarrow ggggh$ sub-amplitude we have,

$$\begin{aligned} H_4^{1234}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}, h) &= \frac{\bar{\mu}^{4-n}}{r_\Gamma} \frac{1}{i\pi^{n/2}} \int d^n \ell \frac{\text{Num}(\ell)}{\prod_i d_i(\ell)} \\ &= \sum_{i,j,k,l} e_{i \times j \times k \times l}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) E_0(p_i, p_j, p_k, p_l; m) \\ &\quad + \sum_{i,j,k} d_{i \times j \times k}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) D_0(p_i, p_j, p_k; m) \\ &\quad + \sum_{i,j} c_{i \times j}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) C_0(p_i, p_j; m) \\ &\quad + \sum_i b_i(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) B_0(p_i; m) + r(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}). \end{aligned} \quad (2.11)$$

The scalar bubble (B_0), triangle (C_0), box (D_0) and pentagon (E_0) integrals, and the constant r_Γ , are defined in appendix A. $\bar{\mu}$ is an arbitrary mass scale, and r are the rational terms. The rank of a Feynman integral is defined to be the number of powers of the loop momentum in the numerator. A scalar Feynman integral has no powers of the loop momentum in the numerator, and is hence of rank zero. All scalar integrals are well known and readily evaluated using existing libraries [18–20]. The sums in the above equation scan over groupings of external gluons. Thus, for example, the sum for the scalar triangle integrals will contain a term $c_{1 \times 234}$ which multiplies the scalar triangle integral $C_0(p_1, p_{234}; m)$ where $p_{234} = p_2 + p_3 + p_4$. The reduction in eq. (2.11) is written in n dimensions, although at the end the amplitude is finite. The individual bubble integrals contain ultra-violet singularities that are regulated using dimensional regularization.

In four dimensions the pentagon integral can be reduced to a sum of the five box integrals obtained by removing one propagator [21–23],

$$\begin{aligned} E_0(p_1, p_2, p_3, p_4; m) = & \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(1)} D_0(p_2, p_3, p_4; m) + \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(2)} D_0(p_{12}, p_3, p_4; m) \\ & + \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} D_0(p_1, p_{23}, p_4; m) \\ & + \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(4)} D_0(p_1, p_2, p_{34}; m) + \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} D_0(p_1, p_2, p_3; m). \end{aligned} \quad (2.12)$$

Explicit forms for the pentagon reduction coefficients, $\mathcal{C}_{1 \times 2 \times 3 \times 4}^{(i)}$, are

$$\begin{aligned} \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(1)} &= -\frac{1}{2} \frac{s_{23} s_{34} [2 s_{12} s_{24} + s_{13} s_{24} + s_{34} s_{12} - s_{23} s_{14}]}{16 |S_{1 \times 2 \times 3 \times 4}|} \\ \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(2)} &= -\frac{1}{2} \frac{s_{34} [s_{1234} s_{23} (s_{123} - 2 s_{12}) + s_{123} (s_{34} (s_{123} - s_{23}) + s_{12} (s_{234} + s_{23}) - s_{234} s_{123})]}{16 |S_{1 \times 2 \times 3 \times 4}|} \\ \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} &= -\frac{1}{2} \frac{[s_{14} s_{23} - (s_{12} + s_{13}) (s_{24} + s_{34})] [s_{34} s_{12} + s_{23} s_{14} - s_{13} s_{24}]}{16 |S_{1 \times 2 \times 3 \times 4}|} \\ \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(4)} &= -\frac{1}{2} \frac{s_{12} [s_{1234} s_{23} (s_{234} - 2 s_{34}) + s_{234} (s_{12} (s_{234} - s_{23}) + s_{34} (s_{123} + s_{23}) - s_{234} s_{123})]}{16 |S_{1 \times 2 \times 3 \times 4}|} \\ \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} &= -\frac{1}{2} \frac{s_{12} s_{23} [2 s_{34} s_{13} + s_{13} s_{24} + s_{34} s_{12} - s_{23} s_{14}]}{16 |S_{1 \times 2 \times 3 \times 4}|} \end{aligned} \quad (2.13)$$

The factor $|S_{1 \times 2 \times 3 \times 4}|$ is the determinant of the matrix, $[S_{1 \times 2 \times 3 \times 4}]_{ij} = [m^2 - \frac{1}{2}(q_{i-1} - q_{j-1})^2]$, where q_i is the offset momentum, see eq. (A.2). It can be written as,

$$\begin{aligned} 16 |S_{1 \times 2 \times 3 \times 4}| &= s_{12} s_{23} s_{34} (s_{14} s_{23} - (s_{12} + s_{13}) (s_{24} + s_{34})) + m^2 G, \\ G &= (s_{12} s_{34} - s_{13} s_{24} - s_{14} s_{23})^2 - 4 s_{13} s_{14} s_{23} s_{24}. \end{aligned} \quad (2.14)$$

As a result of eq. (2.12), in four dimensions the integral basis given in eq. (2.11) is over-complete. The full amplitude can be described by the box, triangle and bubble integrals alone (+ rational terms). This is the specific choice made in this paper, but the other choice to keep the redundant basis of eq. (2.11) is also perfectly viable. In this paper we will work in a basis without pentagon integrals, but the box coefficients will in part display vestiges of their pentagon origin, through effective pentagon coefficients and the presence of the pentagon-to-box reduction coefficients, $\mathcal{C}_{1 \times 2 \times 3 \times 4}^{(i)}$. This will be explained in detail in section 5. Our decomposition of the sub-amplitudes is thus,

$$\begin{aligned} H_4(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) = & \sum_{i,j,k} d_{i \times j \times k} (1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) D_0(p_i, p_j, p_k; m) \\ & + \sum_{i,j} c_{i \times j} (1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) C_0(p_i, p_j; m) \\ & + \sum_i b_i (1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) B_0(p_i; m) + r(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}), \end{aligned} \quad (2.15)$$

which also applies for the $0 \rightarrow \bar{q} q g g h$ sub-amplitude H_4^{34} since it contains no pentagon diagrams in the first place.

2.3 Unitarity methods

The modern treatment of one-loop amplitudes containing massive particles was pioneered 25 years ago in ref. [24]. Since that early paper a whole set of tools and methods have been invented to deal with one-loop amplitudes (for an introduction and comprehensive review see refs. [25] and [26] respectively). We shall apply many of them in order to arrive at the simplest form for the Higgs + 4 parton amplitude. This paper will present compact expressions for the coefficients in the four dimensional version of eq. (2.11) where the scalar pentagon integral has been expressed as a sum of box integrals. The coefficients will be expressed in terms of spinor products. Our notation for spinor products is reported in appendix B.

As we shall see below, there is an intimate connection between the full one-loop calculation with a massive fermion and a suitably normalized calculation performed with the Higgs boson coupling to four partons via a loop of colour-triplet, massive scalar particles. The latter calculation with scalar intermediaries has two advantages. First, the scalar calculation is completely free of Dirac algebra, allowing more compact expressions to be maintained throughout the calculation. This is useful if one can show the identity of the coefficients of scalar integrals between the scalar and the fermionic theories. Second, the scalar calculation, unlike the fermionic calculation, can be performed in the $m \rightarrow 0$ limit. If it can be shown that:

1. the result for a particular coefficient in the scalar theory is identical to the result in the fermionic theory,
2. that particular coefficient is also independent of the mass,

the value of the $m \rightarrow 0$ limit is established.

In order to perform the reduction to scalar integrals indicated in eq. (2.11) we use unitarity techniques to isolate the contribution of boxes [27], triangles [28] and bubbles [29–31]. Since bubble integrals satisfy both criteria enumerated above, their coefficients are most easily calculated using a massless internal scalar loop.

Integrals that do not give rise to rational terms are said to be cut-constructible. In general, x -point integrals are cut constructible in four dimensions if the rank r satisfies $r < \max[(x - 1), 2]$. Thus rank-3 pentagons, rank-2 boxes, rank-1 triangles and bubbles are cut constructible. Integrals that are not cut-constructible give rise to the rational terms (r) in eq. (2.11). In our case the rational terms can instead be obtained by using already-computed results for the mass-dependent coefficients of triangle integrals [32].

2.4 Simplification techniques

We now briefly describe two further techniques that are useful to help simplify the results obtained using unitarity methods. Both methods exploit the fact that the kinematics of our process can be expressed as massless 6-point kinematics by decomposing the momentum of the Higgs boson into two light-like momenta, which for definiteness we call p_5, p_6 .

Reduction of the analytic forms to simpler expressions is aided by the use of momentum twistors [33–36]. In this formalism each particle is described by a 4-component momentum

twistor $Z(\lambda, \mu)$, where λ is the usual two-component holomorphic Weyl spinor (with $\langle i j \rangle = \lambda_\alpha \lambda^\alpha$) and μ is a two-component object related to dual momentum coordinates [33]. Anti-holomorphic spinors ($\tilde{\lambda}_i$, with $[i j] = \tilde{\lambda}^{\dot{\alpha}} \tilde{\lambda}_{\dot{\alpha}}$) are obtained from these via the identity,

$$\tilde{\lambda}_i = \frac{\langle i (i+1) \rangle \mu_{i-1} + \langle (i+1) (i-1) \rangle \mu_i + \langle (i-1) i \rangle \mu_{i+1}}{\langle i (i+1) \rangle \langle (i-1) i \rangle}. \quad (2.16)$$

To describe an n -particle scattering amplitude there are thus $4n$ momentum twistor components, of which only $3n - 10$ are independent due to a $U(1)$ symmetry for each particle and overall Poincaré symmetry. We thus need 8 momentum-twistor variables ($x_1 \dots x_8$) to describe our 6-point kinematics, which we choose to parametrize as,

$$Z = \begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 & \lambda_4 & \lambda_5 & \lambda_6 \\ \mu_1 & \mu_2 & \mu_3 & \mu_4 & \mu_5 & \mu_6 \end{pmatrix} = \begin{pmatrix} 1 & 0 & y_1 & y_2 & y_3 & y_4 \\ 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & x_5 x_6 & x_6 & 1 \\ 0 & 0 & 1 & 1 & 1 - \left(\frac{1-x_7}{x_2 x_5}\right) & -\left(\frac{1+x_8}{x_2 x_5 x_6}\right) \end{pmatrix}, \quad (2.17)$$

where $y_i = \sum_{j=1}^i \prod_{k=1}^j 1/x_k$. The spinors involving our four massless partons are then given by,

$$\begin{aligned} \langle 12 \rangle &= -1, & \langle 13 \rangle &= -1, & \langle 14 \rangle &= -1, \\ \langle 23 \rangle &= 1/x_1, & \langle 24 \rangle &= \frac{1+x_2}{x_1 x_2}, & \langle 34 \rangle &= \frac{1}{x_1 x_2}, \\ [12] &= x_1, & [13] &= x_1 x_8, & [14] &= -\frac{x_1(x_7+x_8)}{x_5}, \\ [23] &= -x_1^2 x_2 x_5 x_6, & [24] &= x_1^2 x_2 x_6, & [34] &= -x_1^2 x_2 x_6 x_7, \end{aligned} \quad (2.18)$$

where the $[i j]$ spinors in the second line have been derived with the aid of eq. (2.16). Note that the variables x_3 and x_4 are not present, leaving us with a rational parametrization of our amplitude in terms of only 6 parameters. Inverting we have

$$\begin{aligned} x_1 &= -\langle 12 \rangle [12], & x_2 &= +\frac{\langle 23 \rangle \langle 14 \rangle}{\langle 12 \rangle \langle 34 \rangle}, & x_3 &= +\frac{\langle 34 \rangle \langle 15 \rangle}{\langle 13 \rangle \langle 45 \rangle}, & x_4 &= -\frac{\langle 45 \rangle \langle 16 \rangle}{\langle 14 \rangle \langle 56 \rangle}, \\ x_5 &= -\frac{\langle 13 \rangle [23]}{\langle 14 \rangle [24]}, & x_6 &= -\frac{\langle 34 \rangle [24]}{\langle 13 \rangle [12]}, & x_7 &= -\frac{\langle 13 \rangle [34]}{\langle 12 \rangle [24]}, & x_8 &= +\frac{\langle 13 \rangle [13]}{\langle 12 \rangle [12]} \end{aligned} \quad (2.19)$$

where we see explicitly that the variables x_3 and x_4 involve momenta p_5 and p_6 and effectively decouple in our case [36]. In order to use this parametrization we first need to remove the overall phase of the coefficient corresponding to the helicities of the external gluons, for example by multiplying by $\langle 12 \rangle^2 \langle 34 \rangle^2$ for the all-plus amplitude. The advantage of this approach is that the amplitude can now be simplified using straightforward algebra, without needing to account for momentum conservation and Schouten identities to manipulate spinor strings. In this way overall factors can easily be identified and the true denominator structure of the coefficients established.

The second method we adopt is to use high precision floating-point arithmetic to simplify our analytic expressions [37]. The study of singular and doubly singular limits in

complex phase space allows us to explore the singularity structure of the integral coefficients. The integral coefficients can then be reconstructed by solving linear systems for the rational numerical coefficients of generic spinor trial functions. The helicities of the gluons impose constraints on the structure of the trial functions. This method is particularly useful when unitarity techniques result in lengthy expressions that are hard to treat using twistor variables, such as in the case of some triangle and bubble coefficients. It is also useful to bypass the algebra involved in removing artefacts of loop-momentum parametrizations, such as square roots and massless projections of non-lightlike external momenta [28, 38]. This paper is the first instance where this method has been applied in the presence of massive particles.

3 Higgs boson production mediated by a coloured scalar

Consider a complex scalar field ϕ in the triplet representation of colour SU(3) coupled to a gluon field and to the Higgs boson, h . The part of the QCD Lagrangian involving the field ϕ is

$$\mathcal{L} = (D_\mu \phi^\dagger)_i (D^\mu \phi)_i - \lambda \phi_i^\dagger \phi_i h \quad (3.1)$$

where $(D_\mu \phi)_i = (\partial_\mu \delta_{ij} + ig(t \cdot \mathcal{A}_\mu)_{ij})\phi_j$. The partial correspondence with the fermion theory emerges when setting $\lambda = -4m^2/v$.

We will calculate colour-ordered sub-amplitudes for the production of a Higgs boson and four gluons mediated by a scalar loop. For the Higgs + 4 gluon amplitude this is,

$$\begin{aligned} \mathcal{A}_4^{gggg}(\{p_i, h_i, c_i\}) = & i \frac{g_s^4}{16\pi^2} \left(-\frac{\lambda}{4} \right) \left\{ [\text{tr}(t^{c_1} t^{c_2} t^{c_3} t^{c_4}) + \text{tr}(t^{c_1} t^{c_4} t^{c_3} t^{c_2})] A_4^{1234}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \right. \\ & + [\text{tr}(t^{c_1} t^{c_3} t^{c_4} t^{c_2}) + \text{tr}(t^{c_1} t^{c_2} t^{c_4} t^{c_3})] A_4^{1342}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \\ & \left. + [\text{tr}(t^{c_1} t^{c_4} t^{c_2} t^{c_3}) + \text{tr}(t^{c_1} t^{c_3} t^{c_2} t^{c_4})] A_4^{1423}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) \right\} \end{aligned} \quad (3.2)$$

and the colour amplitudes have the decomposition,

$$\begin{aligned} A_4^{1234}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}; h) = & \frac{\bar{\mu}^{4-n}}{r_\Gamma} \frac{1}{i\pi^{n/2}} \int d^n \ell \frac{\text{Num}(\ell)}{\prod_i d_i(\ell)} \\ = & \sum_{i,j,k,l} \tilde{e}_{i \times j \times k \times l}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) E_0(p_i, p_j, p_k, p_l; m) \\ & + \sum_{i,j,k} \tilde{d}_{i \times j \times k}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) D_0(p_i, p_j, p_k; m) \\ & + \sum_{i,j} \tilde{c}_{i \times j}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) C_0(p_i, p_j; m) \\ & + \sum_i \tilde{b}_i(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) B_0(p_i; m) + \tilde{r}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}). \end{aligned} \quad (3.3)$$

Thus the tilde indicates that we are referring to an amplitude mediated by a scalar field. For the case of the triangle coefficients we divide the coefficient into two pieces, to separate




Figure 1. Triangle diagrams showing the production of a Higgs boson by gluon fusion.

the mass dependence. Thus for both the fermion- and scalar-mediated loops we have,

$$c_{i \times j} = c_{i \times j}^{(0)} + m^2 c_{i \times j}^{(2)} \quad (3.4)$$

$$\tilde{c}_{i \times j} = \tilde{c}_{i \times j}^{(0)} + m^2 \tilde{c}_{i \times j}^{(2)} \quad (3.5)$$

In eq. (3.2) we have chosen the normalization so that in some cases the coefficients $c_{i \times j} = \tilde{c}_{i \times j}$ and in addition in all cases $c_{i \times j}^{(2)} = \tilde{c}_{i \times j}^{(2)}$, $b_i = \tilde{b}_i$ and $r = \tilde{r}$. Thus we can perform certain parts of the calculations in the scalar theory. We further have that \tilde{b}_i are independent of the mass m , so that they can be calculated in the massless scalar theory. In addition, for the case at hand the rational terms are fully fixed by $\tilde{c}_{i \times j}^{(2)}$,

$$r(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}) = \frac{1}{2} \sum_{i,j} \tilde{c}_{i \times j}^{(2)}(1^{h_1}, 2^{h_2}, 3^{h_3}, 4^{h_4}), \quad (3.6)$$

where the sum runs over all non-zero $\tilde{c}_{i \times j}^{(2)}$ for a particular helicity.

3.1 Relationship of the fermion theory to the scalar theory

In order to elucidate the relationship between the fermion and scalar theories [39] it is instructive to review the steps previously used to demonstrate their similarity using the second order formalism [40]. Following ref. [40] we define the quantity \mathbf{A} to represent the combination of the numerator part of a fermion propagator with momentum ℓ and a gluon-quark-antiquark vertex at which momentum p_1 flows out along the gluon line

$$\mathbf{A}^\mu = (\not{\ell} + m) \gamma^\mu = \left(\not{\ell} + \frac{1}{2} \not{p}_1 - \frac{1}{2} \not{p}_1 + m \right) \gamma^\mu = (2\ell^\mu + p_1^\mu) \mathbf{1} - \frac{1}{2} [\not{p}_1, \gamma^\mu] - \gamma^\mu (\not{\ell} + \not{p}_1 - m). \quad (3.7)$$

The first term on the right hand side of eq. (3.7) already resembles the vertex for a gluon (on-shell or off-shell) interacting with a scalar field. Now consider the integrand of a triangle diagram for a Higgs boson coupled to two gluons as shown in figure (1a),

$$(-1) \times t^{c_1} t^{c_2} \times \frac{1}{m} \frac{\text{tr}\{(\not{\ell} + m) \gamma^{\mu_1} (\not{\ell}_1 + m) \gamma^{\mu_2} (\not{\ell}_{12} + m)\}}{D(\ell) D(\ell_1) D(\ell_{12})} \quad (3.8)$$

where our notation for the loop momenta and propagator factors $D(\ell)$ is given in appendix A. The minus sign is included because of the fermion loop and an overall factor of

$1/m$ has been included for convenience. By using the decomposition in eq. (3.7) twice, the expression in eq. (3.8) can be brought into the form,

$$(-1) \times t^{c_1} t^{c_2} \times \left[\frac{\text{tr}\{\mathbf{B}^{\mu_1}(\ell, \ell_1) \mathbf{B}^{\mu_2}(\ell_1, \ell_{12})\}}{D(\ell) D(\ell_1) D(\ell_{12})} - \frac{\text{tr}\{\gamma^{\mu_1} \gamma^{\mu_2}\}}{D(\ell) D(\ell_{12})} \right], \quad (3.9)$$

where \mathbf{B} is a four-by-four matrix-valued function. It contains a convection term (as expected for a scalar field) and a spin term,

$$\mathbf{B}^\mu(\ell, \ell_1) = (\ell^\mu + \ell_1^\mu) \mathbf{1} + \frac{1}{2}[(\ell - \ell_1), \gamma^\mu] \equiv (\ell^\mu + \ell_1^\mu) \mathbf{1} - \frac{1}{2}[\not{\ell}_1, \gamma^\mu]. \quad (3.10)$$

Note that the spin term is independent of the loop momentum. The result for the companion triangle as shown in figure (1b) is

$$\begin{aligned} & (-1) \times t^{c_2} t^{c_1} \times \frac{1}{m} \frac{\text{tr}\{(-\not{\ell}_{12} + m) \gamma^{\mu_2} (-\not{\ell}_1 + m) \gamma^{\mu_1} (-\not{\ell} + m)\}}{D(\ell) D(\ell_1) D(\ell_{12})} \\ &= (-1) \times t^{c_2} t^{c_1} \times \left[\frac{\text{tr}\{\mathbf{B}^{\mu_2}(-\ell_{12}, -\ell_1) \mathbf{B}^{\mu_1}(-\ell_1, -\ell)\}}{D(\ell) D(\ell_1) D(\ell_{12})} - \frac{\text{tr}\{\gamma^{\mu_2} \gamma^{\mu_1}\}}{D(\ell) D(\ell_{12})} \right]. \end{aligned} \quad (3.11)$$

Adding both diagrams, dropping vanishing terms and exploiting the cyclicity of the trace, the integrand appearing in the full amplitude is,

$$\begin{aligned} & (-1) \left\{ t^{c_1} t^{c_2} \times \left[\frac{(\ell^{\mu_1} + \ell_1^{\mu_1})(\ell_1^{\mu_2} + \ell_{12}^{\mu_2}) \text{tr}\{\mathbf{1}\} + \frac{1}{4} \text{tr}\{\not{\ell}_1, \gamma^{\mu_1}\} [\not{\ell}_2, \gamma^{\mu_2}]}{D(\ell) D(\ell_1) D(\ell_{12})} \right] \right. \\ & \quad + t^{c_2} t^{c_1} \times \left[\frac{(\ell^{\mu_1} + \ell_1^{\mu_1})(\ell_1^{\mu_2} + \ell_{12}^{\mu_2}) \text{tr}\{\mathbf{1}\} + \frac{1}{4} \text{tr}\{\not{\ell}_1, \gamma^{\mu_1}\} [\not{\ell}_2, \gamma^{\mu_2}]}{D(\ell) D(\ell_1) D(\ell_{12})} \right] \\ & \quad \left. - (t^{c_1} t^{c_2} + t^{c_2} t^{c_1}) g^{\mu_1 \mu_2} \frac{\text{tr}\{\mathbf{1}\}}{D(\ell) D(\ell_{12})} \right\}. \end{aligned} \quad (3.12)$$

If we drop the terms involving the commutators of gamma matrices, the fermionic loop (after removing an overall factor of m) can be written as the effect of a (suitably normalized) scalar triangle, with 3-point and 4-point (seagull) vertices. The full fermionic theory requires the inclusion of the additional spin flip terms, given by commutators. These additional terms do not involve the loop momentum and are thus of lower rank. Note also that there is no explicit mass dependence in eq. (3.12). Dependence on the mass m will be generated by the reduction to scalar integrals. In contrast to the full fermionic theory, we may also consider the scalar theory in the massless case.

Iterating this argument for a larger number of gluons, it can be shown that separation of the full fermionic theory into a suitably normalized scalar theory, plus spin terms involving gamma matrix commutators of rank lower by two powers of ℓ , continues to hold. The scalar theory is obtained by dropping all of the spin terms, cf. eq. (3.10). Thus the full amplitude can be written as the sum of the scalar theory and a correction of lower rank, ΔF ,

$$\text{Fermion theory} = \text{Scalar theory} + \Delta F \quad (3.13)$$

The difference between the scalar theory and the full fermionic theory ΔF is of lower rank. Although the scalar theory contains rank-4 pentagons, rank-3 boxes, and rank-2 triangles, ΔF contains only rank-2 pentagons, rank-1 boxes and rank-0 triangles. This has several important consequences:

1. ΔF is cut constructible.
2. ΔF gives no contribution to bubble integral coefficients. Bubble integral coefficients can thus be calculated in the scalar theory.
3. ΔF gives no contribution to the m^2 contributions to triangle coefficients, $c_{i \times j}^{(2)}$.
4. ΔF gives no contribution to certain m^0 triangle coefficients, $c_{i \times j}^{(0)}$.

Exploiting these facts provides the following identities for the $gggh$ case.

$$\begin{aligned} c_{3 \times 4}(1^+, 2^+, 3^+, 4^-) &= \tilde{c}_{3 \times 4}(1^+, 2^+, 3^+, 4^-), \\ c_{2 \times 34}(1^+, 2^+, 3^+, 4^-) &= \tilde{c}_{2 \times 34}(1^+, 2^+, 3^+, 4^-), \\ c_{1 \times 43}(1^+, 2^+, 3^+, 4^-) &= \tilde{c}_{1 \times 43}(1^+, 2^+, 3^+, 4^-), \end{aligned} \quad (3.14)$$

$$\begin{aligned} c_{3 \times 4}(1^+, 2^-, 3^+, 4^-) &= \tilde{c}_{3 \times 4}(1^+, 2^-, 3^+, 4^-), \\ c_{2 \times 34}(1^+, 2^-, 3^+, 4^-) &= \tilde{c}_{2 \times 34}(1^+, 2^-, 3^+, 4^-), \end{aligned} \quad (3.15)$$

$$\begin{aligned} c_{2 \times 3}(1^+, 2^+, 3^-, 4^-) &= \tilde{c}_{2 \times 3}(1^+, 2^+, 3^-, 4^-), \\ c_{1 \times 23}(1^+, 2^+, 3^-, 4^-) &= \tilde{c}_{1 \times 23}(1^+, 2^+, 3^-, 4^-), \end{aligned} \quad (3.16)$$

i.e. all triangles that do not have the Higgs boson as an external leg can be calculated fully in the scalar theory. In appendix C we reproduce several results for tree graphs involving massive scalars, Higgs bosons, gluons and a quark-antiquark pair. These results are useful in applying unitarity to calculate loop diagrams.

4 Coefficients for $H_4^{1234}(g^+, g^+, g^+, g^+; h)$

The reduction of a scalar pentagon integral into a sum of box integrals shown in eq. (2.12) applies in four dimensions. Therefore an extraction of the scalar pentagon integral coefficient must be performed by making use of unitarity methods in d dimensions. To this end the loop momentum is expressed most generally as,

$$\ell^\nu = \alpha p_1^\nu + \beta p_2^\nu + \frac{\gamma}{2} \langle 1 | \gamma^\nu | 2 \rangle + \frac{\delta}{2} \langle 2 | \gamma^\nu | 1 \rangle + \ell_\epsilon^\nu \quad (4.1)$$

where ℓ_ϵ represents the excursion beyond four dimensions, with $\ell_\epsilon^2 = -\mu^2$. Putting the propagators on shell determines α , β , γ , δ and μ^2 . Parametrizing the amplitude at hand with the decomposition in eq. (4.1) we thus find the pentagon coefficient,

$$e_{1 \times 2 \times 3 \times 4}(1^+, 2^+, 3^+, 4^+) = \frac{(m^2 + \mu^2) (s_{1234} - 4(m^2 + \mu^2)) \text{tr}_+ \{1 \, 2 \, 3 \, 4\}}{\langle 1 \, 2 \rangle \langle 2 \, 3 \rangle \langle 3 \, 4 \rangle \langle 4 \, 1 \rangle} \quad (4.2)$$

where the trace functions are defined by,

$$\begin{aligned}\text{tr}_+\{1\,2\,3\,4\} &= \text{tr}\{\gamma_R \not{p}_1 \not{p}_2, \not{p}_3, \not{p}_4\} = [1\,2] \langle 2\,3 \rangle [3\,4] \langle 4\,1 \rangle, \\ \text{tr}_-\{1\,2\,3\,4\} &= \text{tr}\{\gamma_L \not{p}_1 \not{p}_2, \not{p}_3, \not{p}_4\} = \langle 1\,2 \rangle [2\,3] \langle 3\,4 \rangle [4\,1],\end{aligned}\tag{4.3}$$

and $\gamma_{R/L} = (1 \pm \gamma_5)/2$. The identities on the far right of eq. (4.3) holds only for lightlike p_i . The value of μ^2 is fixed by the constraint $\ell^2 = m^2$ (implying $m^2 + \mu^2 = -\gamma\delta s_{12}$) but we have left the coefficient in eq. (4.2) in this form in order to emphasise the d -dimensional nature of the result. We choose to write our amplitude in terms of an effective pentagon coefficient \hat{e} that corresponds to the four-dimensional limit $\mu^2 \rightarrow 0$,

$$\hat{e}_{1 \times 2 \times 3 \times 4}(1^+, 2^+, 3^+, 4^+) = \frac{m^2 (s_{1234} - 4m^2) \text{tr}_+\{1\,2\,3\,4\}}{\langle 1\,2 \rangle \langle 2\,3 \rangle \langle 3\,4 \rangle \langle 4\,1 \rangle}\tag{4.4}$$

This leads to a very compact form for the complete amplitude [15],

$$\begin{aligned}H_4^{1234}(1^+, 2^+, 3^+, 4^+; h) &= \left\{ \frac{4m^2 - s_{1234}}{\langle 1\,2 \rangle \langle 2\,3 \rangle \langle 3\,4 \rangle \langle 4\,1 \rangle} \left[-\text{tr}_+\{1\,2\,3\,4\} m^2 E_0(p_1, p_2, p_3, p_4; m) \right. \right. \\ &\quad + \frac{1}{2}((s_{12} + s_{13})(s_{24} + s_{34}) - s_{14}s_{23}) D_0(p_1, p_{23}, p_4; m) \\ &\quad + \frac{1}{2}s_{12}s_{23} D_0(p_1, p_2, p_3; m) \\ &\quad \left. \left. + (s_{12} + s_{13} + s_{14}) C_0(p_1, p_{234}; m) \right] + 2 \frac{s_{12} + s_{13} + s_{14}}{\langle 1\,2 \rangle \langle 2\,3 \rangle \langle 3\,4 \rangle \langle 4\,1 \rangle} \right\} \\ &\quad + \left\{ 3 \text{ cyclic permutations} \right\}.\end{aligned}\tag{4.5}$$

Although this expression, that includes the scalar pentagon integral, is very simple, we do not follow this approach for the other helicity choices in the following sections. Instead, since in the end our aim is to produce a numerical code to calculate this amplitude, we feel that the structure is more straight-forward working only in terms of boxes, triangles and bubble coefficients. Adopting this approach also for this amplitude means that the minimal set of coefficients that must be specified corresponds to the ones shown in the first and third columns of table 1. Note that coefficients of integrals that could in principle appear but that are not specified in this table (and in subsequent tables in later sections) vanish. There is some ambiguity in the naming convention for the integral coefficients. For a given colour-ordered amplitude the external legs will appear in cyclic or anti-cyclic order, cf. eq. (2.3). For example, $d_{1 \times 2 \times 34}$ in table 1 could equally well be written as $d_{43 \times 2 \times 1}$. Our convention is that the vertex containing more than one gluon, should appear last in the name of the coefficient. Where the compound vertex is in the centre, we have chosen the cyclic ordering.

4.1 Boxes

4.1.1 $d_{1 \times 2 \times 34}$

$$d_{1 \times 2 \times 34}(1^+, 2^+, 3^+, 4^+) = C_{1 \times 2 \times 3 \times 4}^{(4)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^+\}}\tag{4.6}$$

Coefficient	Related coefficients	Coefficient	Related coefficients
$d_{1 \times 2 \times 34}$	$d_{2 \times 3 \times 41}, d_{3 \times 4 \times 12}, d_{4 \times 1 \times 23},$ $d_{1 \times 4 \times 32}, d_{2 \times 1 \times 43}, d_{3 \times 2 \times 14}, d_{4 \times 3 \times 21}$	$c_{1 \times 234}$	$c_{2 \times 341}, c_{3 \times 412}, c_{4 \times 123}$
$d_{1 \times 23 \times 4}$	$d_{2 \times 34 \times 1}, d_{3 \times 41 \times 2}, d_{4 \times 12 \times 3}$		
$d_{1 \times 2 \times 3}$	$d_{2 \times 3 \times 4}, d_{3 \times 4 \times 1}, d_{4 \times 1 \times 2}$		

Table 1. Minimal set of integral coefficients for $1_g^+ 2_g^+ 3_g^+ 4_g^+$.

4.1.2 $d_{1 \times 23 \times 4}$

$$d_{1 \times 23 \times 4}(1^+, 2^+, 3^+, 4^+) = \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^+\}} + \frac{1}{2} \frac{(4m^2 - s_{1234})}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} [(s_{12} + s_{13})(s_{24} + s_{34}) - s_{14} s_{23}] \quad (4.7)$$

4.1.3 $d_{1 \times 2 \times 3}$

$$d_{1 \times 2 \times 3}(1^+, 2^+, 3^+, 4^+) = \mathcal{C}_{4 \times 1 \times 2 \times 3}^{(1)} \hat{e}_{\{4^+ \times 1^+ \times 2^+ \times 3^+\}} + \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^+\}} + \frac{1}{2} \frac{(4m^2 - s_{1234})}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} s_{12} s_{23} \quad (4.8)$$

4.2 Triangles

4.2.1 $c_{1 \times 234}^{(0)}, c_{1 \times 234}^{(2)}$

$$c_{1 \times 234}^{(0)}(1^+, 2^+, 3^+, 4^+) = -(s_{12} + s_{13} + s_{14}) \frac{s_{1234}}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \quad (4.9)$$

$$c_{1 \times 234}^{(2)}(1^+, 2^+, 3^+, 4^+) = 4(s_{12} + s_{13} + s_{14}) \frac{1}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \quad (4.10)$$

In presenting these results we have adopted the notation defined in eq. (3.4) to separately quote the mass-independent and mass-dependent parts.

4.3 Rational terms

$$\begin{aligned} r(1^+, 2^+, 3^+, 4^+) &= \frac{1}{2} [c_{1 \times 234}^{(2)}(1^+, 2^+, 3^+, 4^+) + c_{1 \times 234}^{(2)}(2^+, 3^+, 4^+, 1^+) \\ &\quad + c_{1 \times 234}^{(2)}(3^+, 4^+, 1^+, 2^+) + c_{1 \times 234}^{(2)}(4^+, 1^+, 2^+, 3^+)] \\ &= 4 \frac{s_{1234}}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \end{aligned} \quad (4.11)$$

5 Coefficients for $H_4^{1234}(g^+, g^+, g^+, g^-; h)$

Following the procedure outlined in section 4 to obtain the pentagon coefficients yields, for this particular helicity combination,

$$e_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} = -s_{12} s_{34} (s_{123} - 4(m^2 + \mu^2)) \left[\frac{[23] \langle 4|(2+3)|1]}{\text{tr}_5\{1234\}^2} \right], \quad (5.1)$$

where we have introduced the notation,

$$\text{tr}_5\{1\,2\,3\,4\} = \text{tr}\{\gamma_5 \not{p}_1 \not{p}_2 \not{p}_3 \not{p}_4\} = [1\,2] \langle 2\,3 \rangle [3\,4] \langle 4\,1 \rangle - \langle 1\,2 \rangle [2\,3] \langle 3\,4 \rangle [4\,1] . \quad (5.2)$$

Note that the last identity, eq. (5.2), only applies for the case of lightlike momenta. At this point we could follow the same strategy as in the previous section and take the limit $\mu^2 \rightarrow 0$ to obtain an effective pentagon coefficient \hat{e} . However the appearance of the factor $\text{tr}_5\{1\,2\,3\,4\}$ in the denominator of eq. (5.1) is unpalatable since it is an unphysical singularity. In the amplitude its presence is compensated by corresponding factors in box coefficients, and separating contributions in this way can lead to a loss of numerical precision.

As an alternative solution, we also choose to modify the coefficient itself in such a way that these factors are eliminated. We do so by noting that the denominator can be expressed as,

$$\text{tr}_5\{1\,2\,3\,4\}^2 = (s_{12} s_{34} - s_{14} s_{23} - s_{13} s_{24})^2 - 4 s_{14} s_{23} s_{24} s_{13} = G , \quad (5.3)$$

where G has already been introduced in eq. (2.14). In fact, rearranging that equation and making use of spinor notation we have,

$$s_{12}s_{23}s_{34} \langle 1|(2+3)|4 \rangle \langle 4|(2+3)|1 \rangle = m^2 \text{tr}_5\{1\,2\,3\,4\}^2 - 16 |S_{1 \times 2 \times 3 \times 4}| . \quad (5.4)$$

We may use this equation to eliminate the factor of $\text{tr}_5\{1\,2\,3\,4\}^2$ in the pentagon coefficient indicated in eq. (5.1). The additional term that is introduced contains a factor of $|S_{1 \times 2 \times 3 \times 4}|$; this neatly cancels the denominator factor involved when reducing the pentagon integral to boxes (cf. eq. (2.13)) such that factors of $1/\text{tr}_5\{1\,2\,3\,4\}^2$ can be explicitly absorbed into the box coefficients and cancelled.

In this way we arrive at effective pentagon coefficients,

$$\hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} = (s_{123} - 4m^2) m^2 \left[\frac{[2\,3] \langle 4|(2+3)|1 \rangle}{\langle 2\,3 \rangle \langle 1|(2+3)|4 \rangle} \right] , \quad (5.5)$$

$$\hat{e}_{\{4^- \times 1^+ \times 2^+ \times 3^+\}} = \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} \{1 \leftrightarrow 3\} , \quad (5.6)$$

$$\begin{aligned} \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} = & -m^2 \frac{[2\,3]}{\langle 2\,3 \rangle [3\,4]} \left(\frac{[2\,3] \langle 2|(3+4)|1 \rangle \langle 4|(1+3)|2 \rangle}{\langle 1|(3+4)|2 \rangle} + [1\,3] \langle 4|(2+3)|1 \rangle \right. \\ & \left. + 4m^2 \frac{[2\,3] \langle 3\,4 \rangle \langle 2|(3+4)|1 \rangle}{\langle 2\,3 \rangle \langle 1|(3+4)|2 \rangle} \right) , \end{aligned} \quad (5.7)$$

$$\hat{e}_{\{3^+ \times 4^- \times 1^+ \times 2^+\}} = \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} \{1 \leftrightarrow 3\} . \quad (5.8)$$

The correspondence between $\hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}}$ defined here, and $e_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}}$ given in eq. (5.1) is clear from the first term on the right-hand side of eq. (5.4). We will see that the box coefficients take a particularly simple form when written this way.

The minimal set of remaining integral coefficients to determine this amplitude is shown in the first and third columns of table 2. The related coefficients included in the table are

Coefficient	Related coefficients	Coefficient	Related coefficients
$d_{1 \times 2 \times 34}$	$d_{3 \times 2 \times 14}$	$c_{3 \times 4}$	$c_{4 \times 1}$
$d_{1 \times 4 \times 32}$	$d_{3 \times 4 \times 12}$	$c_{2 \times 34}$	$c_{2 \times 14}$
$d_{2 \times 1 \times 43}$	$d_{2 \times 3 \times 41}$	$c_{1 \times 43}$	$c_{3 \times 41}$
$d_{2 \times 34 \times 1}$	$d_{3 \times 41 \times 2}$	$c_{4 \times 123}$	
$d_{4 \times 3 \times 21}$	$d_{4 \times 1 \times 23}$	$c_{1 \times 234}$	$c_{3 \times 412}$
$d_{1 \times 23 \times 4}$	$d_{4 \times 12 \times 3}$	$c_{2 \times 341}$	
$d_{2 \times 3 \times 4}$	$d_{4 \times 1 \times 2}$	$c_{12 \times 34}$	$c_{23 \times 41}$
$d_{1 \times 2 \times 3}$		b_{34}	b_{14}
$d_{3 \times 4 \times 1}$		b_{234}	b_{412}, b_{341}
		b_{1234}	

Table 2. Minimal set of integral coefficients for $1_g^+ 2_g^+ 3_g^+ 4_g^-$.

determined by using symmetry properties of the amplitude and relabelling momenta. These are mostly straightforward except for the coefficient b_{341} where we have found the relation,

$$b_{341}(1^+, 2^+, 3^+, 4^-) = -b_{234}(2^+, 3^+, 1^+, 4^-) - b_{234}(2^+, 1^+, 3^+, 4^-). \quad (5.9)$$

5.1 Boxes

5.1.1 $d_{1 \times 2 \times 34}$

$$\begin{aligned}
 d_{1 \times 2 \times 34}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(4)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} \\
 &+ \frac{1}{2} \frac{[12][23]\langle 1|(2+4)|3\rangle}{[34]\langle 1|(3+4)|2\rangle\langle 1|(2+3)|4\rangle} \left([23]s_{1234} - 4 \frac{\langle 1|(2+4)|3\rangle}{\langle 12\rangle} m^2 \right) \\
 &- \frac{1}{2} \frac{[12]\langle 24\rangle\langle 4|(2+3)|1\rangle}{\langle 23\rangle\langle 34\rangle\langle 2|(3+4)|1\rangle} \left(\langle 4|(2+3)|1\rangle + 4 \frac{\langle 24\rangle}{\langle 12\rangle} m^2 \right) \quad (5.10)
 \end{aligned}$$

5.1.2 $d_{1 \times 4 \times 32}$

$$\begin{aligned}
 d_{1 \times 4 \times 32}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(2)} \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} \\
 &+ \frac{1}{2} \frac{[23]s_{14}s_{234}}{\langle 23\rangle^2[34]\langle 1|(3+4)|2\rangle\langle 1|(2+3)|4\rangle} (4m^2s_{234} - s_{23}s_{1234}) \quad (5.11)
 \end{aligned}$$

5.1.3 $d_{2 \times 1 \times 43}$

$$\begin{aligned}
 d_{2 \times 1 \times 43}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{3 \times 4 \times 1 \times 2}^{(2)} \hat{e}_{\{3^+ \times 4^- \times 1^+ \times 2^+\}} \\
 &+ \frac{1}{2} \frac{[12]\langle 4|(1+3)|2\rangle^2}{\langle 12\rangle\langle 34\rangle\langle 1|(3+4)|2\rangle\langle 3|(1+4)|2\rangle} (4m^2\langle 14\rangle - \langle 12\rangle\langle 4|(1+3)|2\rangle) \\
 &+ \frac{1}{2} \frac{[12][13]^2}{\langle 12\rangle[14][34]\langle 2|(3+4)|1\rangle} (4m^2\langle 2|(1+4)|3\rangle - \langle 21\rangle[13]s_{1234}) \quad (5.12)
 \end{aligned}$$

5.1.4 $d_{2 \times 34 \times 1}$

$$\begin{aligned}
 d_{2 \times 34 \times 1}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(3)} \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} \\
 &+ \frac{\langle 24 \rangle}{\langle 12 \rangle \langle 23 \rangle} \left[2 \frac{\langle 14 \rangle \langle 24 \rangle \langle 1|(3+4)|2 \rangle \langle 2|(3+4)|1 \rangle}{\langle 12 \rangle^2 \langle 34 \rangle} + \frac{\langle 4|(1+3)|2 \rangle \langle 4|(2+3)|1 \rangle}{2 \langle 34 \rangle} \right. \\
 &\left. + \frac{s_{1234} [13] [23]}{2 [34]} - 2m^2 \left(3 \frac{\langle 14 \rangle \langle 24 \rangle [12]}{\langle 12 \rangle \langle 34 \rangle} + 2 \frac{[13] [23]}{[34]} + \frac{\langle 24 \rangle [14] [23]}{\langle 23 \rangle [34]} \right) \right]
 \end{aligned} \tag{5.13}$$

5.1.5 $d_{4 \times 3 \times 21}$

$$\begin{aligned}
 d_{4 \times 3 \times 21}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(2)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} \\
 &+ (4m^2 - s_{123}) \left(\frac{s_{34} s_{123}^2}{2 \langle 12 \rangle \langle 23 \rangle \langle 1|(2+3)|4 \rangle \langle 3|(1+2)|4 \rangle} \right)
 \end{aligned} \tag{5.14}$$

5.1.6 $d_{1 \times 23 \times 4}$

$$d_{1 \times 23 \times 4}(1^+, 2^+, 3^+, 4^-) = \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} \tag{5.15}$$

5.1.7 $d_{2 \times 3 \times 4}$

$$\begin{aligned}
 d_{2 \times 3 \times 4}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(1)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} + \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(5)} \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} \\
 &+ (4m^2 s_{234} - s_{23} s_{1234}) \frac{\langle 34 \rangle [23]^2}{2 \langle 23 \rangle \langle 1|(3+4)|2 \rangle \langle 1|(2+3)|4 \rangle}
 \end{aligned} \tag{5.16}$$

5.1.8 $d_{1 \times 2 \times 3}$

$$\begin{aligned}
 d_{1 \times 2 \times 3}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} \hat{e}_{\{1^+ \times 2^+ \times 3^+ \times 4^-\}} + \mathcal{C}_{4 \times 1 \times 2 \times 3}^{(1)} \hat{e}_{\{4^- \times 1^+ \times 2^+ \times 3^+\}} \\
 &+ (4m^2 - s_{123}) \frac{s_{123} [12] [23]}{2 \langle 3|(1+2)|4 \rangle \langle 1|(2+3)|4 \rangle}
 \end{aligned} \tag{5.17}$$

Note that, as expected, this whole expression is invariant under $1 \leftrightarrow 3$ since $\mathcal{C}_{4 \times 1 \times 2 \times 3}^{(1)} = \mathcal{C}_{3 \times 2 \times 1 \times 4}^{(5)}$.

5.1.9 $d_{3 \times 4 \times 1}$

$$\begin{aligned}
 d_{3 \times 4 \times 1}(1^+, 2^+, 3^+, 4^-) &= \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(1)} \hat{e}_{\{2^+ \times 3^+ \times 4^- \times 1^+\}} + \mathcal{C}_{3 \times 4 \times 1 \times 2}^{(5)} \hat{e}_{\{3^+ \times 4^- \times 1^+ \times 2^+\}} \\
 &- \frac{1}{2} \frac{s_{14} [23] \langle 34 \rangle}{\langle 13 \rangle \langle 23 \rangle^2 \langle 1|(3+4)|2 \rangle} (\langle 4|(1+3)|2 \rangle \langle 23 \rangle + 4m^2 \langle 34 \rangle) \\
 &- \frac{1}{2} \frac{[12] \langle 14 \rangle s_{34}}{\langle 12 \rangle^2 \langle 13 \rangle \langle 3|(1+4)|2 \rangle} (\langle 4|(1+3)|2 \rangle \langle 21 \rangle + 4m^2 \langle 14 \rangle) \\
 &+ \frac{1}{2} \frac{\langle 14 \rangle \langle 34 \rangle}{\langle 12 \rangle \langle 13 \rangle \langle 23 \rangle} \left[4 \frac{s_{14} s_{34}}{\langle 13 \rangle} + [13] (s_{123} - 12m^2) \right]
 \end{aligned} \tag{5.18}$$

Note that this is manifestly symmetric under the exchange $1 \leftrightarrow 3$.

5.2 Triangles

For the case of the triangle coefficients we divide the coefficient into two pieces, to separate the mass dependence, see eq. (3.4). For many coefficients the $c_{i \times j}^{(2)}$ term is equal to zero. For these cases, the full result is given by the m^0 term and we shall omit the superscript.

5.2.1 $c_{3 \times 4}$

$$c_{3 \times 4}(1^+, 2^+, 3^+, 4^-) = 2s_{34} \frac{\langle 14 \rangle \langle 43 \rangle}{\langle 12 \rangle \langle 23 \rangle \langle 13 \rangle^2}, \quad (5.19)$$

5.2.2 $c_{2 \times 34}$

$$c_{2 \times 34}(1^+, 2^+, 3^+, 4^-) = -2(s_{23} + s_{24}) \frac{\langle 14 \rangle \langle 24 \rangle^2}{\langle 12 \rangle^3 \langle 23 \rangle \langle 34 \rangle} \quad (5.20)$$

5.2.3 $c_{1 \times 43}$

$$c_{1 \times 43}(1^+, 2^+, 3^+, 4^-) = -2(s_{13} + s_{14}) \frac{(\langle 12 \rangle \langle 34 \rangle + \langle 13 \rangle \langle 24 \rangle)}{\langle 12 \rangle \langle 34 \rangle} \left[\frac{\langle 14 \rangle}{\langle 12 \rangle \langle 13 \rangle} \right]^2, \quad (5.21)$$

5.2.4 $c_{4 \times 123}^{(0)}, c_{4 \times 123}^{(2)}$

$$c_{4 \times 123}^{(0)}(1^+, 2^+, 3^+, 4^-) = -\frac{s_{123}^2 (s_{14} + s_{24} + s_{34})}{\langle 12 \rangle \langle 23 \rangle \langle 3|(1+2)|4 \rangle \langle 1|(2+3)|4 \rangle} \quad (5.22)$$

$$c_{4 \times 123}^{(2)}(1^+, 2^+, 3^+, 4^-) = 4 \frac{s_{123} (s_{14} + s_{24} + s_{34})}{\langle 12 \rangle \langle 23 \rangle \langle 3|(1+2)|4 \rangle \langle 1|(2+3)|4 \rangle} \quad (5.23)$$

5.2.5 $c_{1 \times 234}^{(0)}, c_{1 \times 234}^{(2)}$

$$\begin{aligned} c_{1 \times 234}^{(0)}(1^+, 2^+, 3^+, 4^-) = & -2 \frac{\langle 1|(2+3)|1 \rangle^2 \langle 4|(2+3)|1 \rangle}{\langle 1|(2+3)|4 \rangle \langle 12 \rangle \langle 23 \rangle \langle 34 \rangle [14]} \\ & + (s_{12} + s_{13} + s_{14}) \left[\frac{1}{\langle 1|(2+3)|4 \rangle} \left(2 \frac{\langle 4|(2+3)|1 \rangle [23]^2}{s_{34} s_{234}} - \frac{(s_{12} + s_{13})^2 [34] + s_{23} \langle 12 \rangle [23] [14]}{\langle 12 \rangle \langle 13 \rangle \langle 23 \rangle [14] [34]} \right) \right. \\ & \left. + \frac{\langle 3|(1+4)|2 \rangle \langle 14 \rangle [23]^2}{\langle 1|(3+4)|2 \rangle \langle 12 \rangle \langle 13 \rangle [24] s_{34}} - \frac{\langle 4|(2+3)|1 \rangle [13]^2}{\langle 2|(3+4)|1 \rangle \langle 12 \rangle [14] s_{34}} - \frac{\langle 14 \rangle (s_{12}^2 \langle 23 \rangle + 2 \langle 13 \rangle \langle 24 \rangle [14] s_{24})}{\langle 12 \rangle^3 \langle 23 \rangle \langle 13 \rangle \langle 34 \rangle [24] [14]} \right] \end{aligned} \quad (5.24)$$

$$\begin{aligned} c_{1 \times 234}^{(2)}(1^+, 2^+, 3^+, 4^-) = & \frac{8 \langle 4|(2+3)|1 \rangle^3}{s_{234} (s_{12} + s_{13} + s_{14}) \langle 23 \rangle \langle 34 \rangle \langle 2|(3+4)|1 \rangle} \\ & - 4(s_{12} + s_{13} + s_{14}) \left\{ \frac{\langle 14 \rangle^2 [23]}{\langle 12 \rangle^2 \langle 34 \rangle \langle 1|(3+4)|2 \rangle \langle 1|(2+3)|4 \rangle} \right. \\ & - \frac{[23]}{\langle 12 \rangle \langle 1|(2+3)|4 \rangle \langle 1|(3+4)|2 \rangle \langle 2|(3+4)|1 \rangle} \left[\frac{\langle 12 \rangle [13] [23]}{[34]} + \frac{[12] \langle 14 \rangle \langle 24 \rangle}{\langle 34 \rangle} \right] \\ & \left. - \left[\frac{[13] \langle 14 \rangle}{\langle 12 \rangle} + \frac{\langle 4|(2+3)|1 \rangle}{\langle 23 \rangle} \right] \left[\frac{\langle 24 \rangle}{\langle 12 \rangle \langle 34 \rangle \langle 1|(2+3)|4 \rangle \langle 2|(3+4)|1 \rangle} \right] \right\} \end{aligned} \quad (5.25)$$

5.2.6 $c_{2 \times 341}^{(0)}, c_{2 \times 341}^{(2)}$

$$\begin{aligned}
 c_{2 \times 341}^{(0)}(1^+, 2^+, 3^+, 4^-) &= 2 \frac{(s_{12} + s_{23} + s_{24}) \langle 24 \rangle^2 (\langle 14 \rangle^2 \langle 23 \rangle^2 + \langle 12 \rangle^2 \langle 34 \rangle^2)}{\langle 12 \rangle^3 \langle 23 \rangle^3 \langle 14 \rangle \langle 34 \rangle} \\
 &+ \frac{\langle 4|(1+3)|2 \rangle^4}{s_{134} \langle 14 \rangle \langle 34 \rangle \langle 1|(3+4)|2 \rangle \langle 3|(1+4)|2 \rangle} + \frac{(s_{12} + s_{23}) [13]^2}{\langle 12 \rangle [14] \langle 23 \rangle [34]} \\
 &+ \left\{ \frac{(s_{12} + s_{23} + s_{24}) [12] [13]^3}{s_{134} [14] [34] \langle 2|(3+4)|1 \rangle} - \frac{[12] \langle 24 \rangle^2 \langle 4|(3+2)|1 \rangle}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 2|(3+4)|1 \rangle} - \frac{[12] [13]^2 \langle 24 \rangle}{\langle 23 \rangle [34] \langle 2|(3+4)|1 \rangle} \right\} \\
 &+ \{1 \leftrightarrow 3\} \tag{5.26}
 \end{aligned}$$

$$\begin{aligned}
 c_{2 \times 341}^{(2)}(1^+, 2^+, 3^+, 4^-) &= \frac{4(s_{12} + s_{23} + s_{24}) s_{134} [12] [32]}{\langle 12 \rangle \langle 32 \rangle [14] [34] \langle 1|(3+4)|2 \rangle \langle 3|(1+4)|2 \rangle} \\
 &- \frac{8}{(s_{12} + s_{23} + s_{24}) s_{134}} \frac{\langle 4|(1+3)|2 \rangle^4}{\langle 14 \rangle \langle 34 \rangle \langle 1|(3+4)|2 \rangle \langle 3|(1+4)|2 \rangle} \\
 &+ \left\{ \frac{4(s_{12} + s_{23} + s_{24})}{[(s_{13} + s_{14})(s_{23} + s_{24}) - s_{12} s_{34}]} \left[\frac{[13]^2 [23]}{\langle 12 \rangle [14] [34]} - \frac{[12] \langle 14 \rangle \langle 24 \rangle^2}{\langle 12 \rangle^2 \langle 23 \rangle \langle 34 \rangle} \right] \right\} \\
 &+ \{1 \leftrightarrow 3\} \tag{5.27}
 \end{aligned}$$

5.2.7 $c_{12 \times 34}^{(0)}, c_{12 \times 34}^{(2)}$

$$\begin{aligned}
 c_{12 \times 34}^{(0)}(1^+, 2^+, 3^+, 4^-) &= \frac{[13] (s_{12} + s_{23}) \langle 4|(2+3)|1 \rangle}{\langle 12 \rangle \langle 23 \rangle [24] \langle 2|(3+4)|1 \rangle} + \frac{[12] \langle 24 \rangle^2 \langle 4|(2+3)|1 \rangle}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 2|(3+4)|1 \rangle} \\
 &- \frac{[23] \langle 4|(1+3)|2 \rangle^2}{\langle 12 \rangle [24] \langle 34 \rangle \langle 1|(3+4)|2 \rangle} - \frac{[23]^3 s_{1234}}{\langle 12 \rangle [24] [34] \langle 1|(3+4)|2 \rangle} \\
 &- \frac{[13]^2 [23] s_{1234}}{\langle 12 \rangle [24] [34] \langle 2|(3+4)|1 \rangle} + \frac{[12] s_{123} (s_{123} - s_{124})}{\langle 12 \rangle \langle 23 \rangle [24] \langle 3|(1+2)|4 \rangle} \\
 &- \frac{[12] \langle 14 \rangle \langle 4|(2+3)|1 \rangle}{\langle 12 \rangle \langle 23 \rangle [24] \langle 34 \rangle} \tag{5.28}
 \end{aligned}$$

$$\begin{aligned}
 c_{12 \times 34}^{(2)}(1^+, 2^+, 3^+, 4^-) &= -4 \frac{[12]}{\langle 12 \rangle \langle 3|(1+2)|4 \rangle [(s_{13} + s_{14})(s_{23} + s_{24}) - s_{12} s_{34}]} \\
 &\times \left[(\langle 14 \rangle [13] - \langle 24 \rangle [23]) (s_{13} + s_{23} - s_{14} - s_{24}) \right. \\
 &+ \langle 12 \rangle \langle 34 \rangle [13] [23] \left(2 - \frac{(s_{13} + s_{23})(s_{13} + s_{14} + s_{23} + s_{24})}{s_{12} s_{34}} \right) \\
 &\left. + \langle 14 \rangle \langle 24 \rangle [12] [34] \left(2 - \frac{(s_{14} + s_{24})(s_{13} + s_{14} + s_{23} + s_{24})}{s_{12} s_{34}} \right) \right] \tag{5.29}
 \end{aligned}$$

5.3 Bubbles

5.3.1 b_{34}

$$b_{34}(1^+, 2^+, 3^+, 4^-) = \frac{4}{\langle 12 \rangle^2 \langle 13 \rangle \langle 23 \rangle} \left(\frac{\langle 24 \rangle^2 \langle 13 \rangle [23]}{(s_{23} + s_{24})} - \frac{\langle 14 \rangle^2 \langle 23 \rangle [13]}{(s_{13} + s_{14})} \right) \tag{5.30}$$

5.3.2 b_{234}

$$b_{234}(1^+, 2^+, 3^+, 4^-) = \frac{4}{\langle 23 \rangle \langle 34 \rangle} \left(\frac{\langle 24 \rangle^2 \langle 4|(2+3)|1]}{\langle 12 \rangle^2 \langle 2|(3+4)|1]} \right. \\ \left. - \frac{\langle 4|(2+3)|1]^3}{\langle 2|(3+4)|1] (s_{1234} - s_{234})^2} - \frac{\langle 24 \rangle^2 [23] \langle 34 \rangle}{\langle 12 \rangle^2 (s_{23} + s_{24})} \right) \quad (5.31)$$

5.3.3 b_{1234}

Since the full amplitude is finite in four dimensions, one of the coefficients is uniquely determined in terms of the remainder. We thus have,

$$b_{1234}(1^+, 2^+, 3^+, 4^-) = -b_{34} - b_{41} - b_{234} - b_{412} - b_{341}, \quad (5.32)$$

where we have suppressed momentum and helicity labels on the right-hand side for brevity.

5.4 Rational terms

$$r(1^+, 2^+, 3^+, 4^-) = \frac{1}{2} [c_{12 \times 34}^{(2)}(1^+, 2^+, 3^+, 4^-) + c_{12 \times 34}^{(2)}(3^+, 2^+, 1^+, 4^-) \\ + c_{1 \times 234}^{(2)}(1^+, 2^+, 3^+, 4^-) + c_{1 \times 234}^{(2)}(3^+, 2^+, 1^+, 4^-) \\ + c_{4 \times 123}^{(2)}(1^+, 2^+, 3^+, 4^-) + c_{2 \times 341}^{(2)}(1^+, 2^+, 3^+, 4^-)] \quad (5.33)$$

6 Coefficients for $H_4^{1234}(g^+, g^-, g^+, g^-; h)$

For this helicity combination the coefficients of the scalar pentagon integrals contain a factor of $1/\text{tr}_5\{1234\}^4$ and we must modify the pentagon integral coefficients in a similar fashion as described for the $+++-$ configuration in section 5. All coefficients can then be written in terms of a single function,

$$\hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} = -m^2 \frac{\langle 12 \rangle [34] \langle 4|(2+3)|1]}{[12] \langle 34 \rangle \langle 1|(2+3)|4]} \left(\frac{[13]^2 \langle 34 \rangle}{[34]} + \frac{\langle 24 \rangle^2 [12]}{\langle 12 \rangle} + 4m^2 \frac{\langle 4|(2+3)|1]}{\langle 1|(2+3)|4]} \right) \quad (6.1)$$

This is manifestly symmetric under $\{1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow []\}$. Other coefficients are trivially related via symmetries:

$$\hat{e}_{\{3^+ \times 4^- \times 1^+ \times 2^-\}} = \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4 \right\} \quad (6.2)$$

$$\hat{e}_{\{4^- \times 1^+ \times 2^- \times 3^+\}} = \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} \left\{ 1 \rightarrow 4, 2 \rightarrow 1, 3 \rightarrow 2, 4 \rightarrow 3, \langle \rangle \leftrightarrow [] \right\} \quad (6.3)$$

$$\hat{e}_{\{2^- \times 3^+ \times 4^- \times 1^+\}} = \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} \left\{ 1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 4, 4 \rightarrow 1, \langle \rangle \leftrightarrow [] \right\} \quad (6.4)$$

Coefficient	Related coefficients	Coefficient	Related coefficients
$d_{4 \times 3 \times 21}$	$d_{2 \times 1 \times 43}, d_{3 \times 2 \times 14}, d_{1 \times 4 \times 32},$ $d_{1 \times 2 \times 34}, d_{2 \times 3 \times 41},$ $d_{3 \times 4 \times 12}, d_{4 \times 1 \times 23}$	$c_{3 \times 4}$	$c_{4 \times 1}, c_{2 \times 3}, c_{1 \times 2}$
$d_{1 \times 23 \times 4}$	$d_{2 \times 34 \times 1}, d_{3 \times 41 \times 2}, d_{4 \times 12 \times 3}$	$c_{2 \times 34}$	$c_{3 \times 41}, c_{4 \times 12}, c_{1 \times 23}$
$d_{1 \times 2 \times 3}$	$d_{2 \times 3 \times 4}, d_{3 \times 4 \times 1}, d_{4 \times 1 \times 2}$	$c_{12 \times 34}$	$c_{1 \times 43}, c_{2 \times 14}, c_{3 \times 21}, c_{4 \times 32}$
		$c_{1 \times 234}$	$c_{23 \times 41}$
		b_{34}	$c_{2 \times 341}, c_{3 \times 412}, c_{4 \times 123}$
		b_{234}	b_{12}, b_{23}, b_{41}
		b_{1234}	$b_{341}, b_{412}, b_{123}$

Table 3. Minimal set of integral coefficients for $1_g^+ 2_g^- 3_g^+ 4_g^-$.

The minimal set of integral coefficients that must be calculated for the colour ordering H_4^{1234} is shown in the first and third columns of table 3, for example the bubble coefficients are given by:

$$\begin{aligned}
 b_{412}(1^+, 2^-, 3^+, 4^-) &= b_{234}(3^+, 4^-, 1^+, 2^-) \\
 b_{123}(1^+, 2^-, 3^+, 4^-) &= b_{234}(4^+, 1^-, 2^+, 3^-)|_{\langle \rangle \leftrightarrow []} \\
 b_{341}(1^+, 2^-, 3^+, 4^-) &= b_{234}(2^+, 3^-, 4^+, 1^-)|_{\langle \rangle \leftrightarrow []}.
 \end{aligned} \tag{6.5}$$

The calculation of the coefficients of other colour orderings requires the use of $++--$ functions which are given in the next section.

6.1 Boxes

6.1.1 $d_{4 \times 3 \times 21}$

$$\begin{aligned}
 d_{4 \times 3 \times 21}(1^+, 2^-, 3^+, 4^-) &= \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(2)} \\
 &+ \frac{\langle 2|(1+3)|4 \rangle}{\langle 1|(2+3)|4 \rangle \langle 3|(1+2)|4 \rangle} \left[-2 \frac{\langle 23 \rangle \langle 2|(1+3)|4 \rangle s_{34} s_{123}^2}{\langle 12 \rangle \langle 3|(1+2)|4 \rangle^2} \right. \\
 &+ \frac{1}{2} \frac{\langle 24 \rangle^2 [34] s_{123}}{\langle 12 \rangle} + \frac{1}{2} \frac{[13]^2 \langle 34 \rangle s_{123}}{[12]} - 2m^2 \left(2 \frac{[13] \langle 4|(2+3)|1 \rangle}{[12]} \right. \\
 &\left. \left. + \frac{[23] \langle 2|(1+3)|4 \rangle \langle 4|(2+3)|1 \rangle}{[12] \langle 1|(2+3)|4 \rangle} + 3 \frac{\langle 23 \rangle \langle 2|(1+3)|4 \rangle \langle 4|(1+2)|3 \rangle}{\langle 12 \rangle \langle 3|(1+2)|4 \rangle} \right) \right]
 \end{aligned} \tag{6.6}$$

6.1.2 $d_{1 \times 23 \times 4}$

$$\begin{aligned}
 d_{1 \times 23 \times 4}(1^+, 2^-, 3^+, 4^-) &= \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} \\
 &+ \left[\frac{1}{2} \frac{\langle 24 \rangle^3 \langle 4|(2+3)|1 \rangle}{\langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \left(1 + 4m^2 \frac{\langle 12 \rangle}{\langle 24 \rangle \langle 1|(2+3)|4 \rangle} \right) \right] \\
 &+ [1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow []]
 \end{aligned} \tag{6.7}$$

Note that here the symmetrization only applies to the terms in square brackets.

6.1.3 $d_{1 \times 2 \times 3}$

$$\begin{aligned}
 d_{1 \times 2 \times 3}(1^+, 2^-, 3^+, 4^-) &= \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} \hat{e}_{\{1^+ \times 2^- \times 3^+ \times 4^-\}} + \mathcal{C}_{4 \times 1 \times 2 \times 3}^{(1)} \hat{e}_{\{4^- \times 1^+ \times 2^- \times 3^+\}} \\
 &+ \frac{\langle 12 \rangle \langle 23 \rangle}{\langle 1|(2+3)|4 \rangle \langle 3|(1+2)|4 \rangle} \left[-2 \frac{s_{12}s_{23}s_{123}}{\langle 13 \rangle^2} - \frac{1}{2} \frac{[12][23]\langle 24 \rangle^2 (s_{14} + s_{24} + s_{34})}{\langle 14 \rangle \langle 34 \rangle} \right. \\
 &+ \frac{1}{2} [13]^2 s_{123} + 2m^2 \left(2 \frac{[13]s_{123}}{\langle 13 \rangle} - [13]^2 + \frac{[12][23]\langle 24 \rangle^2}{\langle 14 \rangle \langle 34 \rangle} \right. \\
 &\left. \left. - \frac{[23]\langle 2|(1+3)|4 \rangle \langle 4|(2+3)|1 \rangle}{\langle 34 \rangle \langle 1|(2+3)|4 \rangle} + \frac{[12]\langle 2|(1+3)|4 \rangle \langle 4|(1+2)|3 \rangle}{\langle 14 \rangle \langle 3|(1+2)|4 \rangle} \right) \right] \quad (6.8)
 \end{aligned}$$

Note that this is explicitly symmetric under the exchange $1 \leftrightarrow 3$.

6.2 Triangles

6.2.1 $c_{3 \times 4}$

$$\begin{aligned}
 c_{3 \times 4}(1^+, 2^-, 3^+, 4^-) &= \frac{-2s_{34}}{s_{12} \langle 3|(1+2)|4 \rangle^3 \langle 13 \rangle^2 [24]^2} \quad (6.9) \\
 &\left\{ \langle 3|1|4 \rangle^3 [s_{24}(s_{14} + s_{24}) + s_{12}(s_{23} + s_{34})] + \langle 3|1|4 \rangle^2 \langle 3|2|4 \rangle [s_{12}^2 - s_{14}(s_{14} + s_{24}) + s_{12}(3s_{234} - 5s_{24})] \right. \\
 &\left. + \langle 3|2|4 \rangle^3 [s_{13}(s_{13} + s_{23}) + s_{12}(s_{14} + s_{34})] + \langle 3|1|4 \rangle \langle 3|2|4 \rangle^2 [s_{12}^2 - s_{23}(s_{13} + s_{23}) + s_{12}(3s_{134} - 5s_{13})] \right\}
 \end{aligned}$$

6.2.2 $c_{2 \times 34}$

$$c_{2 \times 34}(1^+, 2^-, 3^+, 4^-) = 2(s_{23} + s_{24}) \frac{s_{234} [23]^2 \left(\langle 1|(2+4)|3 \rangle [24] + \langle 1|(3+4)|2 \rangle [34] \right)}{\langle 1|(3+4)|2 \rangle^3 [24]^2 [34]} \quad (6.10)$$

6.2.3 $c_{12 \times 34}^{(0)}, c_{12 \times 34}^{(2)}$

This coefficient is defined in terms of the corresponding coefficient with a scalar loop, $\tilde{c}_{12 \times 34}^{(0)}$

$$\begin{aligned}
 c_{12 \times 34}^{(0)}(1^+, 2^-, 3^+, 4^-) &= \tilde{c}_{12 \times 34}^{(0)}(1^+, 2^-, 3^+, 4^-) \\
 &+ \left\{ \frac{\langle 21 \rangle^2 [13]^2 \langle 34 \rangle^2 - \langle 24 \rangle^2 \langle 1|(3+4)|1 \rangle \langle 4|(1+2)|4 \rangle}{\langle 12 \rangle \langle 34 \rangle \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle} \right\} \quad (6.11) \\
 &+ \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4 \right\} + \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\}
 \end{aligned}$$

$$\begin{aligned}
 c_{12 \times 34}^{(2)}(1^+, 2^-, 3^+, 4^-) &= \left\{ 4 \frac{\langle 2|(3+4)|1 \rangle}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle} \left[\frac{[23]^2 (s_{23} - s_{14})}{[12][34] \langle 1|(3+4)|2 \rangle} + \frac{3}{2} \frac{[13][23]}{[12][34]} \right. \right. \\
 &+ \langle 24 \rangle \frac{(\langle 3|(1+2)|3 \rangle - \langle 4|(1+2)|4 \rangle)}{\Delta_3(1, 2, 3, 4)} \left([23] - \frac{\langle 14 \rangle p_{12} \cdot p_{34}}{\langle 12 \rangle \langle 34 \rangle} \right) \left. \right] \right\} \quad (6.12) \\
 &+ \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4 \right\} + \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\}
 \end{aligned}$$

where Δ_3 is given by eq. (B.2).

The coefficient for a scalar loop is,

$$\begin{aligned}
 \tilde{c}_{12 \times 34}^{(0)}(1^+, 2^-, 3^+, 4^-) = & \left\{ 2 \frac{\langle 23 \rangle^3 [34] \langle 3|(1+2)|3 \rangle (\langle 3|(1+2)|3 \rangle [23] - [12] \langle 14 \rangle [34])}{\langle 12 \rangle \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^3} \right. \\
 & + 2 \frac{\langle 23 \rangle^2 [34] \langle 4|(1+2)|3 \rangle (-2s_{23} - s_{24})}{\langle 12 \rangle \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2} \\
 & + 2 \frac{[12] \langle 23 \rangle^2 [34] (2s_{12}(s_{23} - s_{14} - s_{34}) + 2s_{13}s_{23} + 2s_{23}^2 + s_{14}s_{34} - s_{23}s_{34} + 2[12] \langle 13 \rangle \langle 24 \rangle [34])}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^3} \\
 & + 2 \frac{[14]^2 (\langle 14 \rangle \langle 24 \rangle (2(s_{13} - s_{24}) - 3(s_{34} + s_{14}) - 4(s_{12} + s_{23})) - 2 \langle 13 \rangle [23] \langle 24 \rangle^2 + 3[13] \langle 14 \rangle^2 \langle 23 \rangle)}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2} \\
 & + \frac{s_{14}^2 s_{12} (6s_{13} - 2s_{14} + 2s_{23} + 2s_{24}) - s_{14}^4 + s_{14}^2 s_{23}^2}{\langle 1|(3+4)|2 \rangle^2 \langle 3|(1+2)|4 \rangle^2} - 8 \frac{s_{12} s_{13} s_{14} s_{23}}{\langle 1|(3+4)|2 \rangle^2 \langle 3|(1+2)|4 \rangle^2} \\
 & + 4 \frac{s_{14} s_{1234} \langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)} + 4 \frac{\langle 12 \rangle [13] \langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle \langle 3|(1+4)|2 \rangle}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)} \\
 & + \frac{\langle 1|(2+3)|4 \rangle \langle 2|(3+4)|1 \rangle \langle 3|(1+4)|2 \rangle \langle 4|(1+2)|3 \rangle (\Pi(4, 3, 2, 1) \Pi(1, 2, 3, 4) + \Delta_3(1, 2, 3, 4))}{\langle 1|(3+4)|2 \rangle^2 \langle 3|(1+2)|4 \rangle^2 \Delta_3(1, 2, 3, 4)} \\
 & - 3 \frac{s_{1234} \langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle \Pi(4, 3, 2, 1) \Pi(1, 2, 3, 4) (s_{13} + s_{14} + s_{23} + s_{24})}{2 \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)^2} \\
 & + 5 \frac{s_{1234} \langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle (s_{13} + s_{14} + s_{23} + s_{24})}{2 \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)} - 4 \frac{\langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle} \Big\} \\
 & + \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4 \right\} + \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\} \quad (6.13)
 \end{aligned}$$

where

$$\Pi(i, j, k, l) = s_{ik} + s_{jk} - s_{il} - s_{jl} \quad (6.14)$$

and Δ_3 is given by eq. (B.2).

6.2.4 $c_{1 \times 234}^{(0)}, c_{1 \times 234}^{(2)}$

$$\begin{aligned}
 c_{1 \times 234}^{(0)}(1^+, 2^-, 3^+, 4^-) = & -2 \frac{[13]^4}{[12] [14] [32] [34]} \\
 & + (s_{12} + s_{13} + s_{14}) \left[\frac{[13]^2 [34]}{\langle 1|(2+3)|4 \rangle [14] [23] [24]} + \frac{[13]^2 [32]}{\langle 1|(3+4)|2 \rangle [12] [43] [42]} \right. \\
 & + \frac{\langle 24 \rangle^2 \langle 14 \rangle}{\langle 1|(3+4)|2 \rangle \langle 13 \rangle \langle 12 \rangle \langle 34 \rangle} + \frac{\langle 24 \rangle^2 \langle 12 \rangle}{\langle 1|(2+3)|4 \rangle \langle 13 \rangle \langle 14 \rangle \langle 32 \rangle} \\
 & \left. - 2 \frac{s_{234} \langle 1|(2+4)|3 \rangle^2 \left([23]^2 \langle 1|(2+3)|4 \rangle^2 + [34]^2 \langle 1|(3+4)|2 \rangle^2 \right)}{\langle 1|(2+3)|4 \rangle^3 \langle 1|(3+4)|2 \rangle^3 [23] [34]} \right] \quad (6.15)
 \end{aligned}$$

$$\begin{aligned}
 c_{1 \times 234}^{(2)}(1^+, 2^-, 3^+, 4^-) = & 4 \frac{(s_{12} + s_{13} + s_{14})}{\langle 1|(3+4)|2 \rangle \langle 1|(2+3)|4 \rangle} \left[\left(\frac{\langle 24 \rangle^2}{\langle 23 \rangle \langle 34 \rangle} - \frac{[13]^2}{[12] [14]} \right) \right. \\
 & + \frac{\langle 1|(2+4)|3 \rangle}{\langle 1|(3+4)|2 \rangle} \left(\frac{\langle 14 \rangle \langle 24 \rangle}{\langle 12 \rangle \langle 34 \rangle} + \frac{[13] [23]}{[12] [34]} \right) + \frac{\langle 1|(2+4)|3 \rangle}{\langle 1|(2+3)|4 \rangle} \left(\frac{\langle 12 \rangle \langle 24 \rangle}{\langle 14 \rangle \langle 23 \rangle} + \frac{[13] [34]}{[14] [23]} \right) \\
 & \left. - 8 \frac{[13]^4}{[12] [14] [23] [34] (s_{12} + s_{13} + s_{14})} \right] \quad (6.16)
 \end{aligned}$$

6.3 Bubbles

6.3.1 b_{34}

$$\begin{aligned}
 b_{34}(1^+, 2^-, 3^+, 4^-) = & \left\{ 4 \frac{[1\,3] \langle 1\,4 \rangle^2 s_{134}}{\langle 1\,3 \rangle (s_{13} + s_{14}) \langle 1|(3+4)|2 \rangle^2} \right. \\
 & + 4 \frac{[1\,4]^2 \langle 3\,4 \rangle (s_{13} + s_{14}) \langle 4|(1+2)|3 \rangle (2s_{123} + s_{124})}{[1\,2] \langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2 \Delta_3(1, 2, 3, 4)} \\
 & + 12 \frac{\langle 1\,2 \rangle [1\,4]^2 \langle 3\,4 \rangle \langle 4|(1+2)|3 \rangle s_{134}}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2 \Delta_3(1, 2, 3, 4)} + 4 \frac{\langle 4|(1+2)|3 \rangle \langle 4|(1+3)|4 \rangle}{\langle 1|(3+4)|2 \rangle^2 \langle 3|(1+2)|4 \rangle} \\
 & + 8 \frac{[1\,3] \langle 3\,4 \rangle \langle 1|(2+3)|4 \rangle \langle 4|(1+2)|3 \rangle (s_{234} - s_{134})}{\langle 1|(3+4)|2 \rangle^2 \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)} \\
 & - 2 \frac{[1\,4] \langle 2\,3 \rangle \langle 4|(1+2)|3 \rangle}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2} + 4 \frac{[1\,4] \langle 2\,3 \rangle (s_{13} + s_{14}) \langle 4|(1+2)|3 \rangle (s_{123} - s_{124})}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle^2 \Delta_3(1, 2, 3, 4)} \\
 & + 3 \frac{\langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle (s_{123} - s_{124}) (s_{234} - s_{134}) (s_{134} + s_{234})}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)^2} \\
 & + \frac{\langle 2|(3+4)|1 \rangle \langle 4|(1+2)|3 \rangle (s_{13} - 5s_{14} - 5s_{23} + s_{24} - 14s_{34})}{\langle 1|(3+4)|2 \rangle \langle 3|(1+2)|4 \rangle \Delta_3(1, 2, 3, 4)} \Big\} \\
 & + \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} \tag{6.17}
 \end{aligned}$$

where Δ_3 is given by eq. (B.2).

6.3.2 b_{234}

$$\begin{aligned}
 b_{234}(1^+, 2^-, 3^+, 4^-) = & \left\{ \frac{4s_{234} [3\,4]}{[2\,4]} \left(- \frac{\langle 2\,4 \rangle [3\,4]}{(s_{24} + s_{34}) \langle 1|(2+3)|4 \rangle^2} \right. \right. \\
 & + \frac{[1\,3] \langle 1|(2+4)|3 \rangle}{(s_{1234} - s_{234}) \langle 1|(2+3)|4 \rangle [2\,3] [3\,4]} \left[\frac{[1\,3]}{(s_{1234} - s_{234})} - \frac{[3\,4]}{\langle 1|(2+3)|4 \rangle} \right] \Bigg) \Big\} \\
 & + \left\{ 2 \leftrightarrow 4 \right\} \tag{6.18}
 \end{aligned}$$

6.3.3 b_{1234}

Similarly to eq. (5.32) we have,

$$b_{1234}(1^+, 2^-, 3^+, 4^-) = -b_{34} - b_{12} - b_{23} - b_{41} - b_{234} - b_{341} - b_{412} - b_{123}, \tag{6.19}$$

suppressing momentum and helicity labels on the right-hand side for brevity.

6.4 Rational terms

$$\begin{aligned}
 r(1^+, 2^-, 3^+, 4^-) = & \frac{1}{2} \left[c_{12 \times 34}^{(2)}(1^+, 2^-, 3^+, 4^-) + c_{12 \times 34}^{(2)}(2^+, 3^-, 4^+, 1^-)|_{[\square] \leftrightarrow \langle \rangle} \right. \\
 & + c_{1 \times 234}^{(2)}(1^+, 2^-, 3^+, 4^-) + c_{1 \times 234}^{(2)}(2^+, 3^-, 4^+, 1^-)|_{[\square] \leftrightarrow \langle \rangle} \\
 & \left. + c_{1 \times 234}^{(2)}(3^+, 4^-, 1^+, 2^-) + c_{1 \times 234}^{(2)}(4^+, 1^-, 2^+, 3^-)|_{[\square] \leftrightarrow \langle \rangle} \right] \tag{6.20}
 \end{aligned}$$

Coefficient	Related coefficients	Coefficient	Related coefficients
$d_{1 \times 2 \times 34}$	$d_{2 \times 1 \times 43}, d_{3 \times 4 \times 12}, d_{4 \times 3 \times 21}$	$c_{2 \times 3}$	$c_{4 \times 1}$
$d_{1 \times 4 \times 32}$	$d_{3 \times 2 \times 14}, d_{4 \times 1 \times 23}, d_{2 \times 3 \times 41}$	$c_{1 \times 23}$	$c_{2 \times 14}, c_{3 \times 41}, c_{4 \times 32}$
$d_{2 \times 34 \times 1}$	$d_{4 \times 12 \times 3}$	$c_{23 \times 41}$	
$d_{1 \times 23 \times 4}$	$d_{3 \times 41 \times 2}$	$c_{1 \times 234}$	$c_{2 \times 341}, c_{3 \times 412}, c_{4 \times 123}$
$d_{1 \times 2 \times 3}$	$d_{3 \times 4 \times 1}, d_{4 \times 1 \times 2}, d_{2 \times 3 \times 4}$	b_{23}	b_{41}
		b_{234}	$b_{341}, b_{412}, b_{123}$
		b_{1234}	

Table 4. Minimal set of integral coefficients for $1_g^+ 2_g^+ 3_g^- 4_g^-$.

7 Coefficients for $H_4^{1234}(g^+, g^+, g^-, g^-; h)$

In this case, as for the all-plus helicity amplitude, there are no factors of $1/\text{tr}_5\{1\,2\,3\,4\}^2$ in the pentagon integral coefficients. Therefore the effective pentagon integral coefficients simply correspond to the $\mu^2 \rightarrow 0$ limit, as in the $++++$ case. We thus have,

$$\hat{e}_{\{1^+ \times 2^+ \times 3^- \times 4^-\}} = m^2 (s_{12} + s_{34} - 4m^2) \frac{[1\,2] \langle 3\,4 \rangle}{\langle 1\,2 \rangle [3\,4]} \quad (7.1)$$

$$\hat{e}_{\{2^+ \times 3^- \times 4^- \times 1^+\}} = m^2 \frac{\langle 3\,4 \rangle}{[3\,4]} \left[(s_{34} - 4m^2) \frac{[2\,3] \langle 3\,4 \rangle [4\,1]}{\langle 2\,3 \rangle [3\,4] \langle 4\,1 \rangle} - [1\,2]^2 \right] \quad (7.2)$$

$$\hat{e}_{\{4^- \times 1^+ \times 2^+ \times 3^-\}} = \hat{e}_{\{2^+ \times 3^- \times 4^- \times 1^+\}} \{2 \leftrightarrow 4, 1 \leftrightarrow 3, \langle \rangle \leftrightarrow []\} \quad (7.3)$$

The minimal set of integral coefficients that must be computed in this case is shown in the first and third columns of table 4. For the colour ordering H_4^{1234} the complete set of related coefficients is given in table 4, for example bubble coefficients are given by:

$$\begin{aligned} b_{341}(1^+, 2^+, 3^-, 4^-) &= b_{234}(2^+, 1^+, 4^-, 3^-) \\ b_{123}(1^+, 2^+, 3^-, 4^-) &= b_{234}(4^+, 3^+, 2^-, 1^-)|_{\langle \rangle \leftrightarrow []} \\ b_{412}(1^+, 2^+, 3^-, 4^-) &= b_{234}(3^+, 4^+, 1^-, 2^-)|_{\langle \rangle \leftrightarrow []}. \end{aligned} \quad (7.4)$$

The calculation of the coefficients of other colour orderings requires the use of $+-+--$ functions which are given in the previous section.

7.1 Boxes

7.1.1 $d_{1 \times 2 \times 34}$

$$d_{1 \times 2 \times 34}(1^+, 2^+, 3^-, 4^-) = \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(4)} \hat{e}_{\{1^+ \times 2^+ \times 3^- \times 4^-\}} \quad (7.5)$$

7.1.2 $d_{1 \times 4 \times 32}$

$$\begin{aligned}
 d_{1 \times 4 \times 32}(1^+, 2^+, 3^-, 4^-) = & \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(2)} \hat{e}_{\{2^+ \times 3^- \times 4^- \times 1^+\}} - \frac{[24]}{\langle 1|(2+3)|4\rangle [34]} \\
 & \times \left\{ 2 \frac{[24] s_{14} s_{234}^2 \langle 1|(3+4)|2\rangle}{[23] \langle 1|(2+3)|4\rangle^2} + \frac{1}{2} s_{234} \left[\frac{[14] \langle 34 \rangle^2}{\langle 23 \rangle} + \frac{\langle 14 \rangle [12]^2}{[23]} \right] \right. \\
 & + 2m^2 \left[3 \frac{[24] \langle 1|(3+4)|2\rangle \langle 4|(2+3)|1\rangle}{[23] \langle 1|(2+3)|4\rangle} + \frac{[14] \langle 34 \rangle s_{234}}{\langle 23 \rangle [34]} \right. \\
 & \left. \left. + \frac{\langle 34 \rangle \langle 3|(2+4)|1\rangle}{\langle 23 \rangle} \right] \right\} \quad (7.6)
 \end{aligned}$$

7.1.3 $d_{2 \times 34 \times 1}$

$$\begin{aligned}
 d_{2 \times 34 \times 1}(1^+, 2^+, 3^-, 4^-) = & \mathcal{C}_{2 \times 3 \times 4 \times 1}^{(3)} \hat{e}_{\{2^+ \times 3^- \times 4^- \times 1^+\}} \\
 & + \frac{1}{2} \frac{\langle 34 \rangle \langle 1|(3+4)|2\rangle \langle 2|(3+4)|1\rangle (s_{34} - 4m^2)}{\langle 12 \rangle \langle 14 \rangle \langle 23 \rangle [34]^2} \quad (7.7)
 \end{aligned}$$

7.1.4 $d_{1 \times 23 \times 4}$

$$\begin{aligned}
 d_{1 \times 23 \times 4}(1^+, 2^+, 3^-, 4^-) = & \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(3)} \hat{e}_{\{1^+ \times 2^+ \times 3^- \times 4^-\}} \\
 & - \frac{1}{2} \langle 4|(2+3)|1\rangle \times \left[\frac{[21] [24]}{[14] [23] [34]} \left([21] - \frac{4m^2 [24]}{\langle 1|(2+3)|4\rangle} \right) \right. \\
 & \left. + \frac{\langle 43 \rangle \langle 13 \rangle}{\langle 23 \rangle \langle 14 \rangle \langle 12 \rangle} \left(\langle 43 \rangle - \frac{4m^2 \langle 13 \rangle}{\langle 1|(2+3)|4\rangle} \right) \right] \quad (7.8)
 \end{aligned}$$

7.1.5 $d_{1 \times 2 \times 3}$

$$\begin{aligned}
 d_{1 \times 2 \times 3}(1^+, 2^+, 3^-, 4^-) = & \mathcal{C}_{1 \times 2 \times 3 \times 4}^{(5)} \hat{e}_{\{1^+ \times 2^+ \times 3^- \times 4^-\}} + \mathcal{C}_{4 \times 1 \times 2 \times 3}^{(1)} \hat{e}_{\{4^- \times 1^+ \times 2^+ \times 3^-\}} \\
 & + \frac{[12]^2 \langle 23 \rangle (s_{12} - 4m^2)}{2 [34] \langle 12 \rangle [14]} \quad (7.9)
 \end{aligned}$$

7.2 Triangles

7.2.1 $c_{2 \times 3}$

$$c_{2 \times 3}(1^+, 2^+, 3^-, 4^-) = -2 \frac{s_{23}}{\langle 2|(1+4)|3\rangle^3} \left\{ \frac{[13]^2 s_{134} \langle 2|(3+4)|1\rangle}{[14] [34]} + \frac{\langle 24 \rangle^2 s_{124} \langle 4|(1+2)|3\rangle}{\langle 14 \rangle \langle 12 \rangle} \right\} \quad (7.10)$$

7.2.2 $c_{1 \times 23}$

$$c_{1 \times 23}(1^+, 2^+, 3^-, 4^-) = 2(s_{12} + s_{13}) \frac{s_{123} \langle 13 \rangle^2 \langle 3|(1+2)|4\rangle}{\langle 12 \rangle \langle 23 \rangle \langle 1|(2+3)|4\rangle^3} \quad (7.11)$$

7.2.3 $c_{23 \times 41}^{(0)}, c_{23 \times 41}^{(2)}$

The full coefficient for this scalar integral is defined in terms of the coefficient with a scalar running in the loop, $\tilde{c}_{23 \times 41}^{(0)}$,

$$\begin{aligned} c_{23 \times 41}^{(0)}(1^+, 2^+, 3^-, 4^-) &= \tilde{c}_{23 \times 41}^{(0)}(1^+, 2^+, 3^-, 4^-) \\ &+ \left\{ -\frac{\langle 32 \rangle^2 [21]^2 \langle 14 \rangle^2 - \langle 34 \rangle^2 \langle 2|(1+4)|2 \rangle \langle 4|(2+3)|4 \rangle}{\langle 23 \rangle \langle 14 \rangle \langle 2|(1+4)|3 \rangle \langle 1|(2+3)|4 \rangle} \right\} \\ &+ \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4 \right\} + \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\} \end{aligned} \quad (7.12)$$

In turn the coefficient for a scalar loop is given by,

$$\begin{aligned} \tilde{c}_{23 \times 41}^{(0)}(1^+, 2^+, 3^-, 4^-) &= -\tilde{c}_{12 \times 34}^{(0)}(2^+, 3^-, 1^+, 4^-) \\ &+ \left\{ -2 \Delta(1, 4, 2, 3) \left[\frac{(s_{13} - s_{24})}{\langle 2|(1+4)|3 \rangle \langle 1|(2+3)|4 \rangle} \right]^2 \right. \\ &\quad \left. - 4 \frac{\langle 3|(1+4)|2 \rangle \langle 4|(2+3)|1 \rangle}{\langle 2|(1+4)|3 \rangle \langle 1|(2+3)|4 \rangle} \right\} \end{aligned} \quad (7.13)$$

where $\tilde{c}_{12 \times 34}^{(0)}$ is given by eq. (6.13) with the appropriate permutation of arguments.

$$\begin{aligned} c_{23 \times 41}^{(2)}(1^+, 2^+, 3^-, 4^-) &= \left\{ 4 \frac{\langle 24 \rangle \langle 3|(1+4)|2 \rangle}{\langle 23 \rangle \langle 14 \rangle \langle 2|(1+4)|3 \rangle} \left[\frac{1}{\langle 1|(2+3)|4 \rangle} \left(\frac{\langle 24 \rangle (s_{13} - s_{24})}{\langle 2|(1+4)|3 \rangle} - 2 \langle 34 \rangle \right) \right. \right. \\ &\quad \left. \left. - \frac{\langle 4|(2+3)|1 \rangle}{\Delta_3(1, 4, 2, 3)} \left(\frac{\langle 12 \rangle \langle 3|(1+4)|2 \rangle}{\langle 1|(2+3)|4 \rangle} - \langle 34 \rangle \right) \right] \right\} \\ &+ \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4 \right\} + \left\{ 1 \leftrightarrow 3, 2 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\} \end{aligned} \quad (7.14)$$

where Δ_3 is given by eq. (B.2).

7.2.4 $c_{1 \times 234}^{(0)}, c_{1 \times 234}^{(2)}$

$$\begin{aligned} c_{1 \times 234}^{(0)}(1^+, 2^+, 3^-, 4^-) &= -2(s_{12} + s_{13} + s_{14}) s_{234} \frac{\langle 1|(3+4)|2 \rangle [24]^2}{\langle 1|(2+3)|4 \rangle^3 [23] [34]} + \frac{2 [12]^3}{[14] [23] [34]} \\ &\quad - \frac{(s_{12} + s_{13} + s_{14})}{\langle 12 \rangle \langle 1|(2+3)|4 \rangle [34]} \left(\frac{\langle 13 \rangle \langle 34 \rangle^2 [34]}{\langle 14 \rangle \langle 23 \rangle} - \frac{\langle 12 \rangle [12]^2 [24]}{[14] [23]} \right) \end{aligned} \quad (7.15)$$

$$\begin{aligned} c_{1 \times 234}^{(2)}(1^+, 2^+, 3^-, 4^-) &= \frac{-4}{s_{23} \langle 1|4|3 \rangle} \left\{ \frac{\langle 4|1|2 \rangle^2 (\langle 3|4|1 \rangle - \langle 3|2|1 \rangle)}{s_{14}(s_{12} + s_{13} + s_{14})} \right. \\ &\quad + \frac{[12] \langle 34 \rangle}{\langle 1|(2+3)|4 \rangle} \left(\langle 1|(3+4)|2 \rangle + \frac{(s_{12} + s_{13} + s_{14})}{s_{12}} \langle 1|3|2 \rangle \right) \\ &\quad \left. - 2 \frac{\langle 1|(3+4)|2 \rangle \langle 4|(2+3)|1 \rangle \langle 3|1|2 \rangle}{(s_{12} + s_{13} + s_{14}) \langle 1|(2+3)|4 \rangle} + \frac{\langle 1|(3+4)|2 \rangle^2 \langle 4|(2+3)|1 \rangle \langle 3|1|4 \rangle}{(s_{12} + s_{13} + s_{14}) \langle 1|(2+3)|4 \rangle^2} \right\} \end{aligned} \quad (7.16)$$

7.3 Bubbles

7.3.1 b_{23}

$$\begin{aligned}
 b_{23}(1^+, 2^+, 3^-, 4^-) = & \left\{ -4 \frac{[24]^2 \langle 34 \rangle s_{234}}{[34] \langle 4|(2+3)|4] \langle 1|(2+3)|4]^2} \right. \\
 & -4 \frac{[13]^2 \langle 23 \rangle \langle 1|(2+3)|1] \langle 3|(4+1)|2] (2s_{124} + s_{134} - 3s_{14})}{[14] \langle 1|(2+3)|4] \langle 2|(1+4)|3]^2 \Delta_3(1, 4, 2, 3)} \\
 & + 12 \frac{\langle 14 \rangle [13]^2 \langle 23 \rangle^2 [23] \langle 3|(4+1)|2]}{\langle 1|(2+3)|4] \langle 2|(1+4)|3]^2 \Delta_3(1, 4, 2, 3)} + 4 \frac{\langle 24 \rangle [24] \langle 3|(4+1)|2]}{\langle 1|(2+3)|4]^2 \langle 2|(1+4)|3]} \\
 & - 8 \frac{[24] \langle 23 \rangle \langle 3|(4+1)|2] (\langle 14 \rangle [34] \langle 4|(2+3)|1] + \langle 14 \rangle \langle 23 \rangle [13] [23] + \langle 24 \rangle [23] \langle 4|(2+3)|4])}{\langle 1|(2+3)|4]^2 \langle 2|(1+4)|3] \Delta_3(1, 4, 2, 3)} \\
 & - 4 \frac{\langle 24 \rangle [13] \langle 3|(4+1)|2]}{\langle 1|(2+3)|4] \langle 2|(1+4)|3]^2} + 8 \frac{\langle 14 \rangle [34] [13] \langle 23 \rangle \langle 4|(2+3)|1] \langle 3|(4+1)|2]}{\langle 1|(2+3)|4] \langle 2|(1+4)|3]^2 \Delta_3(1, 4, 2, 3)} \\
 & + 3 \frac{\langle 4|(2+3)|1] \langle 3|(4+1)|2] (s_{124} - s_{134}) (s_{123} - s_{234}) (s_{234} + s_{123})}{\langle 1|(2+3)|4] \langle 2|(1+4)|3] \Delta_3(1, 4, 2, 3)^2} \\
 & + \frac{\langle 4|(2+3)|1] \langle 3|(4+1)|2] (2s_{23} + 5s_{24} + 3s_{34} + 3s_{12} + 5s_{13})}{\langle 1|(2+3)|4] \langle 2|(1+4)|3] \Delta_3(1, 4, 2, 3)} \Big\} \\
 & + \left\{ 1 \leftrightarrow 4, 2 \leftrightarrow 3, \langle \rangle \leftrightarrow [] \right\}
 \end{aligned} \tag{7.17}$$

where Δ_3 is given by eq. (B.2).

7.3.2 b_{234}

$$\begin{aligned}
 b_{234}(1^+, 2^+, 3^-, 4^-) = & \frac{4 s_{234}}{[34] \langle 1|(2+3)|4]^2} \left(\frac{\langle 34 \rangle [24]^2}{(s_{24} + s_{34})} \right. \\
 & \left. + \frac{[12] \langle 1|(3+4)|2]}{[23]} \left[\frac{\langle 1|(2+3)|4] [12]}{(s_{12} + s_{13} + s_{14})^2} - \frac{[24]}{(s_{12} + s_{13} + s_{14})} \right] \right)
 \end{aligned} \tag{7.18}$$

7.3.3 b_{1234}

Similarly to eq. (5.32) we have,

$$b_{1234}(1^+, 2^+, 3^-, 4^-) = -b_{23} - b_{41} - b_{234} - b_{341} - b_{412} - b_{123}, \tag{7.19}$$

suppressing momentum and helicity labels on the right-hand side for brevity.

7.4 Rational terms

$$\begin{aligned}
 r(1^+, 2^+, 3^-, 4^-) = & \frac{1}{2} \left[c_{23 \times 41}^{(2)}(1^+, 2^+, 3^-, 4^-) \right. \\
 & + c_{1 \times 234}^{(2)}(1^+, 2^+, 3^-, 4^-) + c_{1 \times 234}^{(2)}(2^+, 1^+, 4^-, 3^-) \\
 & \left. + c_{1 \times 234}^{(2)}(3^+, 4^+, 1^-, 2^-)|_{[\langle \rangle \leftrightarrow \langle \rangle]} + c_{1 \times 234}^{(2)}(4^+, 3^+, 2^-, 1^-)|_{[\langle \rangle \leftrightarrow \langle \rangle]} \right]
 \end{aligned} \tag{7.20}$$

$1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+$		$1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+$		$1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-$	
Coefficient	Related coefficients	Coefficient	Related coefficients	Coefficient	Related coefficients
$d_{3 \times 21 \times 4}$	$d_{3 \times 4 \times 12}$ $c_{4 \times 12}$	$d_{3 \times 21 \times 4}$	$d_{3 \times 4 \times 12}$ $c_{4 \times 12}$	$c_{4 \times 123}$	$c_{3 \times 412}$ b_{412}
$d_{4 \times 3 \times 21}$		$d_{4 \times 3 \times 21}$		b_{123}	
$c_{3 \times 21}$		$c_{3 \times 21}$			
$c_{12 \times 34}$		$c_{3 \times 4}$			
$c_{4 \times 123}$		$c_{12 \times 34}$			
$c_{3 \times 412}$		$c_{4 \times 123}$			
b_{12}		b_{34}			
b_{123}		b_{12}			
b_{412}		b_{123}			
b_{1234}		b_{1234}			

Table 5. Minimal set of integral coefficients for $H_4^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+)$, $H_4^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+)$ and $H_4^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-)$ together with the related coefficients that can be obtained from the base set.

8 Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^+, g^+; h)$

The coefficients that must be computed for this amplitude are shown in table 5.

8.1 Boxes

8.1.1 $d_{3 \times 21 \times 4}$

$$\begin{aligned}
 d_{3 \times 21 \times 4}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= -2 \frac{\langle 24 \rangle \langle 23 \rangle \text{tr}_- \{ \not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12} \}}{\langle 12 \rangle \langle 34 \rangle^3} \\
 &\quad - \frac{1}{2} \frac{[13] [14] s_{1234}}{[12] \langle 34 \rangle} - \frac{1}{2} \frac{\langle 2|(1+3)|4 \rangle \langle 2|(1+4)|3 \rangle}{\langle 12 \rangle \langle 34 \rangle} \\
 &\quad + 2m^2 \left[\frac{[13] [14]}{[12] \langle 34 \rangle} + 3 \frac{\langle 23 \rangle \langle 24 \rangle [34]}{\langle 12 \rangle \langle 34 \rangle^2} \right] \quad (8.1)
 \end{aligned}$$

where we have introduced the notation, (cf. eqs. (4.3))

$$\text{tr}_- \{ \not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12} \} = \langle 3|(1+2)|4 \rangle \langle 4|(1+2)|3 \rangle = (s_{13} + s_{23})(s_{14} + s_{24}) - s_{12} s_{34} \quad (8.2)$$

8.1.2 $d_{4 \times 3 \times 21}$

$$\begin{aligned}
 d_{4 \times 3 \times 21}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{1}{2} \frac{[34] \langle 2|(1+3)|4 \rangle^2}{\langle 12 \rangle \langle 3|(1+2)|4 \rangle} + \frac{1}{2} \frac{[13]^2 [34] s_{1234}}{[12] \langle 4|(1+2)|3 \rangle} \\
 &\quad + 2m^2 \left[\frac{\langle 23 \rangle [34] \langle 2|(1+3)|4 \rangle}{\langle 12 \rangle \langle 34 \rangle \langle 3|(1+2)|4 \rangle} - \frac{[13] [34] \langle 4|(2+3)|1 \rangle}{[12] \langle 34 \rangle \langle 4|(1+2)|3 \rangle} \right] \quad (8.3)
 \end{aligned}$$

8.2 Triangles

8.2.1 $c_{3 \times 21}$

$$c_{3 \times 21}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) = 2(s_{13} + s_{23}) \frac{\langle 23 \rangle \langle 24 \rangle}{\langle 12 \rangle \langle 34 \rangle^3} \quad (8.4)$$

8.2.2 $c_{12 \times 34}^{(0)}, c_{12 \times 34}^{(2)}$

$$\begin{aligned} c_{12 \times 34}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{[14]^2 (s_{123} - s_{124} - s_{34})}{[12] \langle 34 \rangle \langle 3|(1+2)|4]} - \frac{\langle 23 \rangle^2 [34]^2}{\langle 12 \rangle \langle 34 \rangle \langle 3|(1+2)|4]} \\ &+ \frac{[13]^2 (s_{124} - s_{123} - s_{34})}{[12] \langle 34 \rangle \langle 4|(1+2)|3]} - \frac{\langle 24 \rangle^2 [34]^2}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3]} \\ &- \frac{4[13][14]}{[12] \langle 34 \rangle} \end{aligned} \quad (8.5)$$

$$\begin{aligned} c_{12 \times 34}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{4}{\langle 34 \rangle^2 \langle 3|(1+2)|4]} \left[\frac{[14]^2 \langle 34 \rangle}{[12]} - \frac{\langle 23 \rangle^2 [34]}{\langle 12 \rangle} \right] \\ &+ \frac{4}{\langle 34 \rangle^2 \langle 4|(1+2)|3]} \left[\frac{[13]^2 \langle 34 \rangle}{[12]} - \frac{\langle 24 \rangle^2 [34]}{\langle 12 \rangle} \right] \end{aligned} \quad (8.6)$$

8.2.3 $c_{4 \times 123}^{(0)}, c_{4 \times 123}^{(2)}$

$$\begin{aligned} c_{4 \times 123}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{2 \langle 2|(1+3)|4]^3}{\langle 12 \rangle \langle 23 \rangle \langle 3|(1+2)|4] s_{123}} \\ &- (s_{14} + s_{24} + s_{34}) \left[\frac{\langle 2|(1+4)|3] \langle 2|(1+3)|4]}{\langle 12 \rangle \langle 34 \rangle \text{tr}_{-}\{\not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12}\}} \right. \\ &\left. + \frac{[13][14] s_{1234}}{[12] \langle 34 \rangle \text{tr}_{-}\{\not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12}\}} + \frac{2 \langle 23 \rangle \langle 24 \rangle}{\langle 12 \rangle \langle 34 \rangle^3} \right] \end{aligned} \quad (8.7)$$

$$\begin{aligned} c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{4}{\langle 3|(1+2)|4]} \left\{ \frac{(s_{14} + s_{24} + s_{34})}{\langle 4|(1+2)|3]} \left[\frac{\langle 23 \rangle \langle 24 \rangle [34]}{\langle 12 \rangle \langle 34 \rangle^2} + \frac{[13][14]}{[12] \langle 34 \rangle} \right] \right. \\ &\left. - 2 \frac{\langle 2|(1+3)|4]^3}{\langle 12 \rangle \langle 23 \rangle (s_{14} + s_{24} + s_{34}) s_{123}} \right\} \end{aligned} \quad (8.8)$$

8.2.4 $c_{3 \times 412}^{(0)}, c_{3 \times 412}^{(2)}$

$$\begin{aligned} c_{3 \times 412}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= +2 \frac{\langle 2|(1+4)|3]^2 \langle 1|(2+4)|3]}{\langle 21 \rangle \langle 14 \rangle \langle 4|(1+2)|3] s_{124}} \\ &+ (s_{13} + s_{23} + s_{34}) \left[-2 \frac{\langle 24 \rangle \langle 23 \rangle}{\langle 21 \rangle \langle 43 \rangle^3} \right. \\ &+ \frac{[14][13][43] \langle 43 \rangle}{[21] \langle 43 \rangle \text{tr}_{-}\{\not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12}\}} + \frac{\langle 24 \rangle \langle 23 \rangle [43]^2}{\langle 21 \rangle \langle 43 \rangle \text{tr}_{-}\{\not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12}\}} \\ &\left. - \frac{[14]^2}{[21] \langle 43 \rangle \langle 3|(1+2)|4]} - \frac{[13]^2}{[21] \langle 43 \rangle \langle 4|(1+2)|3]} \right] \end{aligned} \quad (8.9)$$

$$\begin{aligned}
 c_{3 \times 412}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= 4 \frac{(s_{13} + s_{23} + s_{34})}{\text{tr}_{-\{\not{p}_3 \not{p}_{12} \not{p}_4 \not{p}_{12}\}}} \left[\frac{\langle 24 \rangle \langle 23 \rangle [43]}{\langle 21 \rangle \langle 43 \rangle^2} + \frac{[14] [13]}{[21] \langle 43 \rangle} \right] \\
 &\quad - 8 \frac{\langle 2|(1+4)|3 \rangle^2 \langle 1|(2+4)|3 \rangle}{\langle 21 \rangle \langle 14 \rangle \langle 4|(1+2)|3 \rangle (s_{13} + s_{23} + s_{34}) s_{124}} \quad (8.10)
 \end{aligned}$$

8.3 Bubbles

8.3.1 b_{12}

$$b_{12}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) = \frac{4}{\langle 34 \rangle^2} \left[\frac{[13] \langle 23 \rangle}{(s_{13} + s_{23})} - \frac{[14] \langle 24 \rangle}{(s_{14} + s_{24})} \right] \quad (8.11)$$

8.3.2 b_{123}

$$\begin{aligned}
 b_{123}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle^2} \left[\frac{\langle 34 \rangle \langle 24 \rangle \langle 2|(1+3)|4 \rangle^2}{(s_{14} + s_{24} + s_{34})^2} \right. \\
 &\quad \left. - \frac{\langle 23 \rangle \langle 24 \rangle \langle 2|(1+3)|4 \rangle}{(s_{14} + s_{24} + s_{34})} - \frac{\langle 12 \rangle [13] \langle 23 \rangle^2}{(s_{13} + s_{23})} \right] \quad (8.12)
 \end{aligned}$$

8.3.3 b_{412}

$$\begin{aligned}
 b_{412}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= 4 \frac{1}{\langle 12 \rangle \langle 14 \rangle \langle 34 \rangle^2} \left[\frac{\langle 23 \rangle \langle 34 \rangle \langle 2|(1+4)|3 \rangle \langle 1|(2+4)|3 \rangle}{(s_{13} + s_{23} + s_{34})^2} \right. \\
 &\quad \left. + \frac{\langle 13 \rangle \langle 24 \rangle \langle 2|(1+4)|3 \rangle}{(s_{13} + s_{23} + s_{34})} - \frac{\langle 12 \rangle \langle 24 \rangle s_{14}}{(s_{14} + s_{24})} \right] \quad (8.13)
 \end{aligned}$$

8.3.4 b_{1234}

The final bubble coefficient is given by the relation,

$$b_{1234}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) = -b_{12} - b_{123} - b_{412}, \quad (8.14)$$

where momentum and helicity labels have been suppressed on the right-hand side.

8.4 Rational terms

$$\begin{aligned}
 r(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) &= \frac{1}{2} \left[c_{12 \times 34}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) + c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) \right. \\
 &\quad \left. + c_{3 \times 412}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+) \right] \quad (8.15)
 \end{aligned}$$

9 Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^-, g^+; h)$

The coefficients that must be computed for this amplitude are shown in table 5.

9.1 Boxes

9.1.1 $d_{3 \times 21 \times 4}$

$$\begin{aligned}
 d_{3 \times 21 \times 4}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) &= \frac{1}{2} \langle 3|(1+2)|4 \rangle \left[\frac{[14]^2}{[21] [43]} - \frac{\langle 23 \rangle^2}{\langle 12 \rangle \langle 34 \rangle} \right] \\
 &\quad + 2m^2 \frac{\langle 3|(1+2)|4 \rangle}{\langle 4|(1+2)|3 \rangle} \left[\frac{\langle 24 \rangle \langle 23 \rangle}{\langle 12 \rangle \langle 34 \rangle} - \frac{[13] [14]}{[12] [34]} \right] \quad (9.1)
 \end{aligned}$$

9.1.2 $d_{4 \times 3 \times 21}$

$$d_{4 \times 3 \times 21}^{(0)}(1_q^+, 2_q^-, 3_g^-, 4_g^+) = \frac{1}{2} \frac{s_{123}}{\langle 4|(1+2)|3]} \left[\frac{\langle 23 \rangle^2 [34]}{\langle 12 \rangle} + \frac{[14]^2 \langle 34 \rangle}{[12]} - 4 \frac{[13] \langle 4|(2+3)|1]}{[12] \langle 4|(1+2)|3]^2} s_{34} s_{123} \right] \\ - 2 \frac{m^2}{\langle 4|(1+2)|3]} \left[\frac{3 [13] \langle 3|(1+2)|4] \langle 4|(2+3)|1]}{[12] \langle 4|(1+2)|3]} + \frac{\langle 23 \rangle \langle 2|(1+3)|4]}{\langle 12 \rangle} \right] \quad (9.2)$$

9.2 Triangles

9.2.1 $c_{3 \times 4}$

$$c_{3 \times 4}(1_q^+, 2_q^-, 3_g^-, 4_g^+) = \left\{ \frac{2 s_{34} [13] \langle 23 \rangle}{\langle 4|(1+2)|3]^2} + \frac{2 s_{34} [13]^2 \langle 34 \rangle (2 s_{12} + s_{13} + s_{23})}{[12] \langle 4|(1+2)|3]^3} \right\} \\ - \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} \quad (9.3)$$

9.2.2 $c_{3 \times 21}$

$$c_{3 \times 21}(1_q^+, 2_q^-, 3_g^-, 4_g^+) = 2 (s_{13} + s_{23}) s_{123} \frac{[13] \langle 4|(2+3)|1]}{[12] \langle 4|(1+2)|3]^3} \quad (9.4)$$

9.2.3 $c_{12 \times 34}^{(0)}, c_{12 \times 34}^{(2)}$

$$c_{12 \times 34}^{(0)}(1_q^+, 2_q^-, 3_g^-, 4_g^+) = 8 (s_{124} - s_{123}) (s_{12} + s_{34} + 2 s_{13} + 2 s_{23}) \frac{\langle 24 \rangle [13] \langle 3|(1+2)|4]}{\langle 4|(1+2)|3]^2 \Delta_3(1, 2, 3, 4)} \\ + \left((9 s_{13} - 7 s_{23} - s_{14} - s_{24} + 4 s_{34}) \langle 24 \rangle [14] \right. \\ \left. - (9 s_{14} - 7 s_{24} - s_{13} - s_{23} + 4 s_{34}) \langle 23 \rangle [13] \right) \times \frac{1}{\langle 4|(1+2)|3]^2} \\ + 12 \frac{s_{1234} ((s_{13} + s_{23})^2 - (s_{14} + s_{24})^2) \langle 2|(3+4)|1] \langle 3|(1+2)|4]}{\langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)^2} \\ + 4 \left\{ 3 (s_{12} + s_{34}) + 4 (s_{13} + s_{23} + s_{14}) \right\} [13] \langle 23 \rangle \\ - \left\{ 3 (s_{12} + s_{34}) + 4 (s_{13} + s_{24} + s_{14}) \right\} [14] \langle 24 \rangle \right) \times \frac{\langle 3|(1+2)|4]}{\langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)} \\ - 24 \frac{[13] \langle 24 \rangle \langle 3|(1+2)|4]^2}{\langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)} - 8 \frac{[14] \langle 23 \rangle \langle 3|(1+2)|4]}{\Delta_3(1, 2, 3, 4)} + 8 \frac{[14] \langle 23 \rangle}{\langle 4|(1+2)|3]} \\ + \left\{ \frac{2 \langle 24 \rangle^2 [34] (s_{14} + s_{24})^2}{\langle 12 \rangle \langle 4|(1+2)|3]^3} + \frac{[13] \langle 24 \rangle (s_{14} + s_{24}) (4 s_{124} - 2 s_{34})}{\langle 4|(1+2)|3]^3} \right. \\ \left. + \frac{(s_{13} + s_{23}) \langle 23 \rangle \langle 24 \rangle (s_{14} + s_{24} - s_{13} - s_{23})}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3]^2} + \frac{2 \langle 23 \rangle \langle 24 \rangle [34] (s_{14} + s_{24})}{\langle 12 \rangle \langle 4|(1+2)|3]^2} \right\} \\ - \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\} \quad (9.5)$$

$$\begin{aligned}
 c_{12 \times 34}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = & \left\{ \frac{4 \langle 23 \rangle^2}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3]} - \frac{4 \langle 24 \rangle^2 \langle 3|(1+2)|4]}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3]^2} \right. \\
 & - 8(s_{13} + s_{23} + s_{14} + s_{24}) \frac{\langle 23 \rangle \langle 24 \rangle \langle 3|(1+2)|4]}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)} \\
 & - 16 \frac{[13] \langle 23 \rangle \langle 3|(1+2)|4]}{\langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)} \left. \right\} \\
 & - \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\}
 \end{aligned} \tag{9.6}$$

9.2.4 $c_{4 \times 123}^{(0)}, c_{4 \times 123}^{(2)}$

$$\begin{aligned}
 c_{4 \times 123}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = & - \frac{2(s_{14} + s_{24} + s_{34}) [13] \langle 4|(2+3)|1] s_{123}}{[12] \langle 4|(1+2)|3]^3} \\
 & - \frac{\langle 23 \rangle^2 (s_{14} + s_{24} + s_{34})}{\langle 12 \rangle \langle 34 \rangle \langle 4|(1+2)|3]} - \frac{[14]^2 (s_{14} + s_{24} + s_{34})}{[12] [34] \langle 4|(1+2)|3]} + \frac{2 [14]^2 [24]}{[12] [23] [34]}
 \end{aligned} \tag{9.7}$$

$$\begin{aligned}
 c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = & \frac{4(s_{14} + s_{24} + s_{34})}{\langle 4|(1+2)|3]^2} \left[\frac{\langle 23 \rangle \langle 24 \rangle}{\langle 12 \rangle \langle 34 \rangle} + \frac{[13] [14]}{[12] [34]} \right] \\
 & - 8 \frac{[14]^2 [24]}{[12] [23] [34] (s_{14} + s_{24} + s_{34})}
 \end{aligned} \tag{9.8}$$

9.3 Bubbles

9.3.1 b_{34}

$$\begin{aligned}
 b_{34}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = & 2 \frac{\langle 3|(1+2)|4]}{s_{12} \langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)} \\
 & \times \left[3 \frac{\langle 2|(3+4)|1] (s_{124}^2 - s_{123}^2) (s_{13} + s_{14} + s_{23} + s_{24})}{\Delta_3(1, 2, 3, 4)} \right. \\
 & + 2 \frac{(s_{124} - s_{123}) (\langle 12 \rangle [13] \langle 4|(2+3)|1] - \langle 24 \rangle [12] \langle 2|(1+4)|3])}{\langle 4|(1+2)|3]} \\
 & \left. - 3(s_{123} + s_{124}) (\langle 23 \rangle [13] - \langle 24 \rangle [14]) \right]
 \end{aligned} \tag{9.9}$$

9.3.2 b_{12}

$$\begin{aligned}
 b_{12}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = & \left\{ \frac{4}{\langle 4|(1+2)|3]^2} \left[- \frac{\langle 12 \rangle [13]^2 \langle 3|(1+2)|4]}{[34] (s_{13} + s_{23})} \right. \right. \\
 & + \frac{\langle 3|(1+2)|4]}{\langle 34 \rangle \Delta_3(1, 2, 3, 4)} \left[s_{12} \langle 24 \rangle [13] \langle 34 \rangle - 2 \langle 12 \rangle [13]^2 \langle 34 \rangle^2 \right. \\
 & \left. \left. - \langle 23 \rangle \langle 4|(1+2)|3] \langle 4|(2+3)|1] - \langle 24 \rangle^2 [12] (s_{13} + s_{23} + s_{14} + s_{24}) \right] \right] \\
 & - 12(s_{13} + s_{23}) (2s_{12} + s_{13} + s_{23}) \frac{\langle 2|(3+4)|1] \langle 3|(1+2)|4]}{\langle 4|(1+2)|3] \Delta_3(1, 2, 3, 4)^2} \left. \right\} \\
 & - \left\{ 1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow [] \right\}
 \end{aligned} \tag{9.10}$$

9.3.3 b_{123}

$$b_{123}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = 4 s_{123} \left[\frac{[1\,4]^2 \langle 4|(1+3)|2\rangle}{[1\,2] [2\,3] \langle 4|(1+2)|3\rangle (s_{14} + s_{24} + s_{34})^2} + \frac{[1\,4] \langle 4|(2+3)|1\rangle}{[1\,2] \langle 4|(1+2)|3\rangle^2 (s_{14} + s_{24} + s_{34})} + \frac{\langle 2\,3\rangle [1\,3]}{(s_{13} + s_{23}) \langle 4|(1+2)|3\rangle^2} \right] \quad (9.11)$$

9.3.4 b_{1234}

The final bubble coefficient is given by the relation,

$$b_{1234}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = -b_{12} - b_{34} - b_{123} - b_{412}, \quad (9.12)$$

where momentum and helicity labels have been suppressed on the right-hand side.

9.4 Rational terms

$$r(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) = \frac{1}{2} [c_{12 \times 34}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) + c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+) - c_{4 \times 123}^{(2)}(2_{\bar{q}}^+, 1_q^-, 4_g^-, 3_g^+)|_{[\leftrightarrow \langle \rangle]}] \quad (9.13)$$

10 Coefficients for $H_4^{34}(\bar{q}^+, q^-, g^+, g^-; h)$

Most of the coefficients for this amplitude can be easily obtained from those for $H_4^{34}(\bar{q}^+, q^-, g^-, g^+)$ by performing the following operation:

$$1 \leftrightarrow 2, \quad \langle \rangle \leftrightarrow []. \quad (10.1)$$

However, for some coefficients, this procedure effectively changes the colour ordering of the gluons in the sub-amplitude. For this reason it is necessary to specify the four coefficients shown in table 5. Results for the base set of coefficients are given explicitly here.

10.1 Triangle

10.1.1 $c_{4 \times 123}^{(0)}, c_{4 \times 123}^{(2)}$

$$c_{4 \times 123}^{(0)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-) = -2 \frac{\langle 3\,4\rangle \langle 2|(1+3)|4\rangle^2 s_{123}}{\langle 1\,2\rangle \langle 3|(1+2)|4\rangle^3} + 2 \frac{\langle 2\,4\rangle \langle 2|(1+3)|4\rangle s_{123}}{\langle 1\,2\rangle \langle 3|(1+2)|4\rangle^2} - \frac{\langle 2\,4\rangle^2 \langle 2|(1+3)|4\rangle}{\langle 1\,2\rangle \langle 2\,3\rangle \langle 3|(1+2)|4\rangle} + \frac{[1\,3]^2 (s_{14} + s_{24} + s_{34})}{[1\,2] [3\,4] \langle 3|(1+2)|4\rangle} - \frac{\langle 2\,4\rangle^3}{\langle 1\,2\rangle \langle 2\,3\rangle \langle 3\,4\rangle} \quad (10.2)$$

$$c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-) = -4 \frac{(s_{14} + s_{24} + s_{34})}{\langle 3|(1+2)|4\rangle^2} \left[\frac{\langle 2\,3\rangle \langle 2\,4\rangle}{\langle 1\,2\rangle \langle 3\,4\rangle} + \frac{[1\,3] [1\,4]}{[1\,2] [3\,4]} \right] + 8 \frac{\langle 2\,4\rangle^3}{\langle 1\,2\rangle \langle 2\,3\rangle \langle 3\,4\rangle (s_{14} + s_{24} + s_{34})} \quad (10.3)$$

10.2 Bubble

10.2.1 b_{123}

$$b_{123}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-) = 4 s_{123} \left[- \frac{\langle 24 \rangle^2 \langle 2|(1+3)|4]}{\langle 12 \rangle \langle 23 \rangle \langle 3|(1+2)|4] (s_{14} + s_{24} + s_{34})^2} \right. \\ \left. - \frac{\langle 24 \rangle \langle 2|(1+3)|4]}{\langle 12 \rangle \langle 3|(1+2)|4]^2 (s_{14} + s_{24} + s_{34})} + \frac{[13] \langle 23 \rangle}{(s_{13} + s_{23}) \langle 3|(1+2)|4]^2} \right] \quad (10.4)$$

10.3 Rational terms

$$r(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-) = \frac{1}{2} \left[c_{12 \times 34}^{(2)}(2_{\bar{q}}^+, 1_q^-, 3_g^-, 4_g^+) |_{[\leftrightarrow \langle \rangle]} + c_{4 \times 123}^{(2)}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-) \right. \\ \left. - c_{4 \times 123}^{(2)}(2_{\bar{q}}^+, 1_q^-, 4_g^+, 3_g^-) |_{[\leftrightarrow \langle \rangle]} \right] \quad (10.5)$$

11 Amplitude for $0 \rightarrow \bar{q}q\bar{q}qh$

The amplitude for the ggh process with off-shell gluons with momenta k_1, k_2 has been given in ref. [9]. Thus our result is exactly given by ref. [9] and is only included here for completeness.

The amplitude can be obtained by considering the tensor current for $0 \rightarrow ggh$ with two off-shell gluons (with momenta k_1 and k_2),

$$\mathcal{T}^{\mu_1 \mu_2}(k_1, k_2) = -i \delta^{c_1 c_2} \frac{g_s^2}{8\pi^2} \left(\frac{m^2}{v} \right) \left[F_T(k_1, k_2) T_T^{\mu_1 \mu_2} + F_L(k_1, k_2) T_L^{\mu_1 \mu_2} \right]. \quad (11.1)$$

The two tensor structures appearing here are,

$$T_T^{\mu_1 \mu_2} = k_1 \cdot k_2 g^{\mu_1 \mu_2} - k_1^{\mu_2} k_2^{\mu_1} \quad (11.2)$$

$$T_L^{\mu_1 \mu_2} = k_1^2 k_2^2 g^{\mu_1 \mu_2} - k_1^2 k_2^{\mu_1} k_2^{\mu_2} - k_2^2 k_1^{\mu_1} k_1^{\mu_2} + k_1 \cdot k_2 k_1^{\mu_1} k_2^{\mu_2} \quad (11.3)$$

and the form factors are given by¹

$$F_T(k_1, k_2) = -\frac{1}{\Delta(k_1, k_2)} \left\{ k_{12}^2 (B_0(k_1; m) + B_0(k_2; m) - 2B_0(k_{12}; m) - 2k_1 \cdot k_2 C_0(k_1, k_2; m)) \right. \\ \left. + (k_1^2 - k_2^2) (B_0(k_1; m) - B_0(k_2; m)) \right\} - k_1 \cdot k_2 F_L(k_1, k_2) \quad (11.4)$$

$$F_L(k_1, k_2) = -\frac{1}{\Delta(k_1, k_2)} \left\{ \left[2 - \frac{3k_1^2 k_2 \cdot k_{12}}{\Delta(k_1, k_2)} \right] (B_0(k_1; m) - B_0(k_{12}; m)) \right. \\ \left. + \left[2 - \frac{3k_2^2 k_1 \cdot k_{12}}{\Delta(k_1, k_2)} \right] (B_0(k_2; m) - B_0(k_{12}; m)) \right. \\ \left. - \left[4m^2 + k_1^2 + k_2^2 + k_{12}^2 - 3 \frac{k_1^2 k_2^2 k_{12}^2}{\Delta(k_1, k_2)} \right] C_0(k_1, k_2; m) - 2 \right\} \quad (11.5)$$

¹Note that to produce our standard overall normalization for the helicity amplitude we have changed the normalization of the form factors by a factor of 2 with respect to ref. [9].

where $k_{12} = k_1 + k_2$ and $\Delta(k_1, k_2) = k_1^2 k_2^2 - (k_1 \cdot k_2)^2$. By contracting eq. (11.3) with currents for the quark-antiquark lines we then arrive at the result for the amplitude. All helicity combinations can be obtained from permutations of the single expression,

$$H_4^{4q}(1_{\bar{q}}^+, 2_q^-, 3_{\bar{q}'}^+, 4_{q'}^-; h) = \left[\frac{\langle 2|(3+4)|1\rangle \langle 4|(1+2)|3\rangle + \langle 24\rangle [13] (2p_{12} \cdot p_{34})}{s_{12} s_{34}} \right] F_T(p_{12}, p_{34}) \\ + 2 \langle 24\rangle [13] F_L(p_{12}, p_{34}). \quad (11.6)$$

12 Large mass limit

In the limit of large (top quark) mass our results agree with those obtained in the effective field theory given by eq. (1.1). For $ggggh$ we have [41–43],

$$m^2 H_4^{1234}(1^+, 2^+, 3^+, 4^+; h) \rightarrow \frac{2}{3} \frac{s_{1234}^2}{\langle 12\rangle \langle 23\rangle \langle 34\rangle \langle 41\rangle} \quad (12.1)$$

$$m^2 H_4^{1234}(1^+, 2^+, 3^+, 4^-; h) \rightarrow \frac{2}{3} \left[-\frac{s_{1234}^2 [13]^4}{s_{134} [43] [41] \langle 2|(1+4)|3\rangle \langle 2|(3+4)|1\rangle} \right. \\ \left. + \frac{\langle 4|(2+3)|1\rangle^3}{s_{234} \langle 2|(3+4)|1\rangle \langle 34\rangle \langle 23\rangle} + \frac{\langle 4|(2+1)|3\rangle^3}{s_{124} \langle 2|(1+4)|3\rangle \langle 14\rangle \langle 21\rangle} \right] \quad (12.2)$$

$$m^2 H_4^{1234}(1^+, 2^+, 3^-, 4^-; h) \rightarrow \frac{2}{3} \left[\frac{[12]^4}{[12] [23] [34] [41]} + \frac{\langle 34\rangle^4}{\langle 12\rangle \langle 23\rangle \langle 34\rangle \langle 41\rangle} \right]. \quad (12.3)$$

In this limit the amplitudes satisfy the dual Ward identity. For example,

$$H_4^{1234}(1^+, 2^-, 3^+, 4^-; h) + H_4^{1234}(3^+, 1^+, 2^-, 4^-; h) + H_4^{1234}(1^+, 3^+, 2^-, 4^-; h) = 0, \quad (12.4)$$

which means that in this limit our amplitude $H_4^{1234}(1^+, 2^-, 3^+, 4^-; h)$ is related to two amplitudes of the form already specified in eq. (12.3). We note in passing that the dual Ward identity means that, in the effective theory, the subleading-color term represented by the second line of eq. (2.4) is absent.

For the large mass limit of the $\bar{q}qggh$ amplitudes we have [42],

$$m^2 H^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^+; h) \rightarrow \frac{2}{3} \left[\frac{\langle 2|(1+4)|3\rangle^2 [14]}{s_{124} \langle 24\rangle} \left[\frac{1}{s_{12}} + \frac{1}{s_{14}} \right] \right. \\ \left. - \frac{\langle 2|(1+3)|4\rangle^2 [13]}{s_{123} s_{12} \langle 23\rangle} + \frac{\langle 2|(3+4)|1\rangle^2}{[12] \langle 23\rangle \langle 24\rangle \langle 34\rangle} \right] \quad (12.5)$$

$$m^2 H^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^+, 4_g^-; h) \rightarrow \frac{2}{3} \left[\frac{\langle 24\rangle^3}{\langle 12\rangle \langle 23\rangle \langle 34\rangle} - \frac{[13]^3}{[12] [14] [34]} \right] \quad (12.6)$$

$$m^2 H^{34}(1_{\bar{q}}^+, 2_q^-, 3_g^-, 4_g^+; h) \rightarrow \frac{2}{3} \left[\frac{\langle 23\rangle^2 \langle 13\rangle}{\langle 12\rangle \langle 14\rangle \langle 34\rangle} - \frac{[14]^2 [24]}{[12] [23] [34]} \right] \quad (12.7)$$

and for $\bar{q}q\bar{q}'q'$,

$$m^2 H^{4q}(1_{\bar{q}}^+, 2_q^-, 3_{\bar{q}'}^+, 4_{q'}^-; h) \rightarrow -\frac{2}{3} \left[\frac{[13]^2}{[12] [34]} + \frac{\langle 24\rangle^2}{\langle 12\rangle \langle 34\rangle} \right]. \quad (12.8)$$

13 Conclusions

We have presented analytic results for all helicity amplitudes representing the processes $0 \rightarrow ggggh$, $0 \rightarrow \bar{q}qgggh$ and $0 \rightarrow \bar{q}q\bar{q}'q'h$, where the interaction is mediated by a loop of massive fermions and all dependence on the fermion mass is retained. In order to obtain compact results we have used unitarity techniques and also exploited the correspondence between this amplitude and the one in which the massive fermion is replaced by a coloured scalar. In order to further simplify our analytic results to the forms presented here we have supplemented this approach by the use of momentum twistors and reconstruction from high-precision numerical evaluations. In combination this powerful set of tools rendered this calculation tractable.

Our results for the amplitudes were checked using an in-house implementation of the D -dimensional unitarity method [44] and also against a previous unitarity-based calculation [10]. Complete agreement was found at the amplitude level. Our results for the squared matrix elements are also in full agreement with those obtained using the code OpenLoops 2 [14]. A comparison of the evaluation time of squared matrix elements against both the previous code implemented in MCFM [45–47] and OpenLoops 2 indicates a speed-up by at least an order of magnitude over previously-available results. Our results have been implemented in version 9.1 of the code MCFM,² that includes a calculation of the full matrix elements for the three partonic processes under consideration. Results are given in the subdirectory `MCFM-9.1/src/ggHgg_mass` of that release and the result for a particular equation can be found by searching for the text “Implementation”; every file that gives the result for an integral coefficient contains a comment of the form:

```
!      Implementation of eq.~(x.xx) from arXiv:2002.04018 v2
```

The results of this paper will be useful for improving calculations of the h +jet process at NLO in the full theory. The analytic forms presented here will also be useful in their own right, for understanding the structure of gauge-theory amplitudes. In particular the simplification of our results due to the choice of effective pentagon integral coefficients may have deeper origins that remain to be explored.

Acknowledgments

We would like to acknowledge useful discussions with Simon Badger and thank Daniel Maître for the use of his code to generate spinor helicity ansätze. This document was prepared using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359.

²The code can be downloaded from <https://mcfm.fnal.gov>.

A Integrals

We define the denominators of the integrals as follows,

$$D(\ell) = \ell^2 - m^2 + i\varepsilon. \quad (\text{A.1})$$

The momenta running through the propagators are,

$$\begin{aligned} \ell_1 &= \ell + p_1 = \ell + q_1 \\ \ell_{12} &= \ell + p_1 + p_2 = \ell + q_2 \\ \ell_{123} &= \ell + p_1 + p_2 + p_3 = \ell + q_3 \\ \ell_{1234} &= \ell + p_1 + p_2 + p_3 + p_4 = \ell + q_4. \end{aligned} \quad (\text{A.2})$$

The p_i are the external momenta, whereas the q_i are the off-set momenta in the propagators. In terms of these denominators the integrals are,

$$\begin{aligned} B_0(p_1; m) &= \frac{\bar{\mu}^{4-n}}{r_\Gamma} \frac{1}{i\pi^{n/2}} \int d^n \ell \frac{1}{D(\ell) D(\ell_1)}, \\ C_0(p_1, p_2; m) &= \frac{1}{i\pi^2} \int d^4 \ell \frac{1}{D(\ell) D(\ell_1) D(\ell_{12})}, \\ D(\ell)(p_1, p_2, p_3; m) &= \frac{1}{i\pi^2} \int d^4 \ell \frac{1}{D(\ell) D(\ell_1) D(\ell_{12}) D(\ell_{123})}, \\ E_0(p_1, p_2, p_3, p_4; m) &= \frac{1}{i\pi^2} \int d^4 \ell \frac{1}{D(\ell) D(\ell_1) D(\ell_{12}) D(\ell_{123}) D(\ell_{1234})}, \end{aligned} \quad (\text{A.3})$$

where $r_\Gamma = 1/\Gamma(1-\epsilon) + O(\epsilon^3)$ and $\bar{\mu}$ is an arbitrary mass scale.

B Spinor algebra

All results are presented using the standard notation for the kinematic invariants of the process,

$$s_{ij} = (p_i + p_j)^2, s_{ijk} = (p_i + p_j + p_k)^2, s_{ijkl} = (p_i + p_j + p_k + p_l)^2, \quad (\text{B.1})$$

and the Gram determinant,

$$\Delta_3(i, j, k, l) = (s_{ijkl} - s_{ij} - s_{kl})^2 - 4s_{ij}s_{kl}. \quad (\text{B.2})$$

We express the amplitudes in terms of spinor products defined as,

$$\langle i j \rangle = \bar{u}_-(p_i) u_+(p_j), \quad [i j] = \bar{u}_+(p_i) u_-(p_j), \quad \langle i j \rangle [j i] = 2p_i \cdot p_j, \quad (\text{B.3})$$

and we further define the spinor sandwiches for massless momenta j and k ,

$$\begin{aligned} \langle i | (j + k) | l \rangle &= \langle i j \rangle [j l] + \langle i k \rangle [k l] \\ [i | (j + k) | l \rangle &= [i j] \langle j l \rangle + [i k] \langle k l \rangle \end{aligned} \quad (\text{B.4})$$

In the Weyl representation for the Dirac gamma matrices we have,

$$\not{p} = \gamma^0 p_0 - \gamma^1 p_1 - \gamma^2 p_2 - \gamma^3 p_3 = \begin{pmatrix} 0 & 0 & p^+ & p^1 - ip^2 \\ 0 & 0 & p^1 + ip^2 & p^- \\ p^- & -p^1 + ip^2 & 0 & 0 \\ -p^1 - ip^2 & p^+ & 0 & 0 \end{pmatrix}, \quad (\text{B.5})$$

where $p^\pm = p^0 \pm p^3$. The spinor solutions of the massless Dirac equation are,

$$u_+(p) = \begin{bmatrix} \sqrt{p^+} \\ \sqrt{p^-} e^{i\varphi_p} \\ 0 \\ 0 \end{bmatrix}, \quad u_-(p) = \begin{bmatrix} 0 \\ 0 \\ \sqrt{p^-} e^{-i\varphi_p} \\ -\sqrt{p^+} \end{bmatrix}, \quad (\text{B.6})$$

where

$$e^{\pm i\varphi_p} \equiv \frac{p^1 \pm ip^2}{\sqrt{(p^1)^2 + (p^2)^2}} = \frac{p^1 \pm ip^2}{\sqrt{p^+ p^-}}, \quad p^\pm = p^0 \pm p^3. \quad (\text{B.7})$$

In this representation the Dirac conjugate spinors are,

$$\bar{u}_+(p) \equiv u_+^\dagger(p) \gamma^0 = [0, 0, \sqrt{p^+}, \sqrt{p^-} e^{-i\varphi_p}] \quad (\text{B.8})$$

$$\bar{u}_-(p) \equiv u_-^\dagger(p) \gamma^0 = [\sqrt{p^-} e^{i\varphi_p}, -\sqrt{p^+}, 0, 0] \quad (\text{B.9})$$

C Results for tree-level amplitudes with massive scalars

In the following we shall give results for colour-ordered tree amplitudes for a scalar, anti-scalar pair coupled to n partons (n gluons or $n - 2$ gluons with a quark antiquark pair). For the n gluon case the amplitudes are defined as follows,

$$\mathcal{A}_n^{\text{tree}}(\ell; 1, \dots, n; \bar{\ell}) = i g_s^n \sum_{\sigma \in \Sigma_n} (t^{c_{\sigma(1)}} \dots t^{c_{\sigma(n)}})_{j\bar{j}} A_n^{\text{tree}}(\ell; \sigma(1), \dots, \sigma(n); \bar{\ell}), \quad (\text{C.1})$$

where Σ_n is the permutation group on n elements, and A_n^{tree} are the tree-level partial amplitudes. The t matrices are the SU(3) matrices in the fundamental representation normalized as in eq. (2.2). The massive scalars are in the fundamental representation of SU(3) and the colour indices of the scalar and anti-scalar are j and \bar{j} respectively.

We adopt throughout the convention that all momenta are taken to be outgoing. Correspondingly we have the tree amplitudes with an additional Higgs boson, derived using the Lagrangian, eq. (3.1),

$$\mathcal{A}_n^{\text{tree}}(h; \ell; 1, \dots, n; \bar{\ell}) = i \left(\frac{-\lambda}{4} \right) g_s^n \sum_{\sigma \in \Sigma_n} (t^{a_{\sigma(1)}} \dots t^{a_{\sigma(n)}})_{j\bar{j}} A_n^{\text{tree}}(h; \sigma(1), \dots, \sigma(n)). \quad (\text{C.2})$$

We define the denominators of the scalar propagators as $D(\ell) = \ell^2 - m^2$, which must be supplemented by the $+i\varepsilon$ prescription when the propagators are used in loop diagrams, cf. eq. (A.1). The momenta in the propagators are defined in eq. (A.2), with similar expressions for $\bar{\ell}$. Because the external momentum p_1 is light-like, for an on-shell ℓ we may also write $D(\ell_1) = \langle 1|\ell|1 \rangle$.

C.1 One gluon

$$A_1^{\text{tree}}(\ell; 1^+; \bar{\ell}) = \frac{\langle b|\ell|1\rangle}{\langle b\,1\rangle}, \quad A_1^{\text{tree}}(\ell; 1^-; \bar{\ell}) = \frac{\langle 1|\ell|b\rangle}{[1\,b]} \quad (\text{C.3})$$

$$\begin{aligned} A_1^{\text{tree}}(h, \ell; 1^+; \bar{\ell}) &= -4 \left[\frac{\langle b|\bar{\ell}|1\rangle}{\langle b\,1\rangle \langle 1|\bar{\ell}|1\rangle} - \frac{\langle b|\ell|1\rangle}{\langle b\,1\rangle \langle 1|\ell|1\rangle} \right] \\ &= -4 \left[\frac{\langle b|\bar{\ell}|1\rangle \langle 1|\ell|1\rangle - \langle b|\ell|1\rangle \langle 1|\bar{\ell}|1\rangle}{\langle b\,1\rangle \langle 1|\bar{\ell}|1\rangle \langle 1|\ell|1\rangle} \right] \end{aligned} \quad (\text{C.4})$$

In eqs. (C.3), (C.4) b is an arbitrary light-like momentum.

C.2 Two gluons

$$A_2^{\text{tree}}(\ell; 1^+, 2^+; \bar{\ell}) = -\frac{m^2 [1\,2]}{\langle 1\,2\rangle D(\ell_1)} \quad (\text{C.5})$$

$$A_2^{\text{tree}}(\ell; 1^+, 2^-; \bar{\ell}) = -\frac{\langle 2|\ell|1\rangle^2}{\langle 1|2|1\rangle D(\ell_1)} \quad (\text{C.6})$$

$$A_2^{\text{tree}}(h, \ell; 1^+, 2^+; \bar{\ell}) = -4 \frac{[1\,2]}{\langle 1\,2\rangle} \left\{ \left[\frac{m^2}{D(\ell_1) D(\ell_{12})} + \frac{m^2}{D(\bar{\ell}_2) D(\bar{\ell}_{12})} \right] - \frac{1}{s_{12}} \left[1 - \frac{\langle 1|\bar{\ell}|2\rangle \langle 2|\ell|1\rangle}{D(\ell_1) D(\bar{\ell}_2)} \right] \right\} \quad (\text{C.7})$$

$$A_2^{\text{tree}}(h, \ell; 1^+, 2^-; \bar{\ell}) = -4 \frac{1}{s_{12}} \left\{ \frac{\langle 2|\ell|1\rangle^2}{D(\ell_{12}) D(\ell_1)} + \frac{\langle 2|\bar{\ell}|1\rangle^2}{D(\bar{\ell}_{12}) D(\bar{\ell}_2)} - \frac{\langle 2|\bar{\ell}|1\rangle \langle 2|\ell|1\rangle}{D(\ell_1) D(\bar{\ell}_2)} \right\} \quad (\text{C.8})$$

C.3 Three gluons

The spurious poles in the original BCFW form of the amplitudes [48] can be eliminated and the amplitudes rewritten in the following form [49],

$$A_3^{\text{tree}}(\ell; 1^+, 2^+, 3^-; \bar{\ell}) = \frac{\langle 3|\bar{\ell}(1+2)\ell|1\rangle \langle 3|\bar{\ell}|2\rangle}{\langle 1\,2\rangle s_{23} D(\ell_1) D(\ell_{12})} + \frac{[1\,2] \langle 3|(1+2)|\bar{\ell}|3\rangle}{\langle 1\,2\rangle s_{23} s_{123}}, \quad (\text{C.9})$$

$$\begin{aligned} A_3^{\text{tree}}(\ell; 1^+, 2^-, 3^+; \bar{\ell}) &= \frac{1}{s_{12} s_{23}} \left[\frac{[2\,1] \langle 2|\ell|1\rangle \langle 2|\bar{\ell}|3\rangle^2}{D(\ell) D(\ell_{12})} + \frac{[3\,1] \langle 2|\ell|1\rangle \langle 2|\bar{\ell}|3\rangle}{D(\ell_{12})} \right. \\ &\quad \left. - \frac{\langle 2|\bar{\ell}|(1+3)|2\rangle [1\,3]^2}{s_{123}} \right]. \end{aligned} \quad (\text{C.10})$$

By using charge conjugation on eq. (C.9) we also obtain the following result,

$$A_3^{\text{tree}}(\ell; 1^+, 2^-, 3^-; \bar{\ell}) = \frac{\langle 3|\bar{\ell}(2+3)\ell|1\rangle \langle 2|\ell|1\rangle}{[2\,3] s_{12} D(\ell_1) D(\ell_{12})} - \frac{\langle 2\,3\rangle [1|(2+3)|\ell|1\rangle}{[2\,3] s_{12} s_{123}}, \quad (\text{C.11})$$

C.4 One gluon, two quarks

For the calculation of the $H^{34}(\bar{q}qgg; h)$ we also need the tree amplitude for a pair of SU(3) triplet scalars coupled to a quark-antiquark pair with an additional gluon.

$$\begin{aligned} \mathcal{A}_3^{\bar{q}qg}(\ell; 1_{\bar{q}}^+, 2_q^-, 3_g^+; \bar{\ell}) = i\sqrt{2}g_s^3 \left\{ \left[(t^{c3})_{jj_1} \delta_{j_2, \bar{j}} - (t^{c3})_{j_2 \bar{j}} \delta_{j, j_1} \right] \frac{[1\,3] \langle 2|\ell| (1+2+3) |2\rangle}{\langle 2\,3\rangle s_{12} s_{123}} \right. \\ - \left[(t^{c3})_{jj_1} \delta_{j_2, \bar{j}} - \frac{1}{N} (t^{c3})_{j_2 j_1} \delta_{j, \bar{j}} \right] \frac{\langle 2|\ell| (1+2+3) |2\rangle}{\langle 1\,3\rangle \langle 2\,3\rangle s_{123}} \\ + \left[(t^{c3})_{jj_1} \delta_{j_2, \bar{j}} - \frac{1}{N} (t^{c3})_{j\bar{j}} \delta_{j_2, j_1} \right] \frac{\langle 2|\ell_{123}|1\rangle \langle 2|\ell|3\rangle}{\langle 2\,3\rangle s_{12} D(\ell_3)} \\ \left. + \left[(t^{c3})_{j_2 \bar{j}} \delta_{j, j_1} - \frac{1}{N} (t^{c3})_{j\bar{j}} \delta_{j_2, j_1} \right] \frac{\langle 2|\ell|1\rangle \langle 2|\ell_{123}|3\rangle}{\langle 2\,3\rangle s_{12} D(\ell_{12})} \right\} \quad (C.12) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_3^{\bar{q}qg}(\ell; 1_{\bar{q}}^+, 2_q^-, 3_g^-; \bar{\ell}) = i\sqrt{2}g_s^3 \left\{ - \left[(t^{c3})_{jj_1} \delta_{j_2, \bar{j}} - (t^{c3})_{j_2 \bar{j}} \delta_{j, j_1} \right] \frac{\langle 2\,3\rangle [1|\ell| (1+2+3) |1\rangle}{[1\,3] s_{12} s_{123}} \right. \\ - \left[(t^{c3})_{j_2 \bar{j}} \delta_{j, j_1} - \frac{1}{N} (t^{c3})_{j_2 j_1} \delta_{j, \bar{j}} \right] \frac{[1|\ell| (1+2+3) |1\rangle}{[1\,3] [2\,3] s_{123}} \\ - \left[(t^{c3})_{jj_1} \delta_{j_2, \bar{j}} - \frac{1}{N} (t^{c3})_{j\bar{j}} \delta_{j_2, j_1} \right] \frac{\langle 2|\ell_{123}|1\rangle [1|\ell|3\rangle}{[1\,3] s_{12} D(\ell_3)} \\ \left. - \left[(t^{c3})_{j_2 \bar{j}} \delta_{j, j_1} - \frac{1}{N} (t^{c3})_{j\bar{j}} \delta_{j_2, j_1} \right] \frac{\langle 2|\ell|1\rangle [1|\ell_{123}|3\rangle}{[1\,3] s_{12} D(\ell_{12})} \right\} \quad (C.13) \end{aligned}$$

D Numerical value of coefficients at a given phase-space point

In order to assist in the reconstruction of the coefficients in a numerical program we give the value of all the needed base set of integral coefficients at a given data point. The point we choose is given by (with $p = (E, p_x, p_y, p_z)$),

$$\begin{aligned} p_1 &= (-15\kappa, -10\kappa, +11\kappa, +2\kappa) \\ p_2 &= (-9\kappa, +8\kappa, +1\kappa, -4\kappa) \\ p_3 &= (-21\kappa, +4\kappa, -13\kappa, +16\kappa) \\ p_4 &= (-7\kappa, +2\kappa, -6\kappa, +3\kappa) \\ p_h &= (+52\kappa, -4\kappa, +7\kappa, -17\kappa) \end{aligned} \quad (D.1)$$

with $\kappa = 1/\sqrt{94} \text{ GeV}$ and $p_h = -p_1 - p_2 - p_3 - p_4$. This fixes $s_{1234} = 25 \text{ GeV}^2$, $M_h = 5 \text{ GeV}$ and we further choose $m = 1.5 \text{ GeV}$. The numerical values of the coefficients and the rational terms are given in tables 6, 7 and 8.

Helicities	Coefficient	Real Part	Imaginary Part	Absolute Value
++++	$d_{1 \times 2 \times 34}$	1.7494424584	0.9145464014	1.9740678903
	$d_{1 \times 23 \times 4}$	5.9642590946	8.6101701300	10.4741308095
	$d_{1 \times 2 \times 3}$	11.9903841330	27.4958119676	29.9964829174
	$c_{1 \times 234}$	18.9100021625	56.3253050259	59.4148817052
	r	-6.4366316747	-19.1721417745	20.2237792595
+++-	$d_{1 \times 2 \times 34}$	-24.3908884307	-34.1026538098	41.9273948071
	$d_{1 \times 4 \times 32}$	-22.2037730441	-29.7427434881	37.1165505885
	$d_{2 \times 1 \times 43}$	16.2217246906	-62.9923572563	65.0475320412
	$d_{2 \times 34 \times 1}$	-66.4392574700	12.9335349956	67.6864185834
	$d_{4 \times 3 \times 21}$	8.2313626631	-0.7960661671	8.2697673869
	$d_{1 \times 23 \times 4}$	2.0815256682	-1.7746340633	2.7353382179
	$d_{2 \times 3 \times 4}$	-0.9920798783	-1.5084323993	1.8054336843
	$d_{1 \times 2 \times 3}$	22.2417370205	1.7066361068	22.3071170816
	$d_{3 \times 4 \times 1}$	0.8741489856	-5.3830902459	5.4536040418
	$c_{3 \times 4}$	-0.0041638038	0.0115576710	0.0122848289
	$c_{2 \times 34}$	3.1035163815	-0.1080335333	3.1053961381
	$c_{1 \times 43}$	6.9656648763	-0.8139894264	7.0130639492
	$c_{4 \times 123}$	12.7875856866	2.0711796271	12.9542322327
	$c_{1 \times 234}$	-41.8343835373	-39.3169799861	57.4102827129
	$c_{2 \times 341}$	-0.0578594858	-18.9964204402	18.9965085545
	$c_{12 \times 34}$	12.4596639704	-35.5399553316	37.6607441672
	b_{34}	-0.0409808246	0.1015477837	0.1095051613
	b_{234}	0.2936947594	-0.0490382211	0.2977605730
	b_{1234}	-0.9341272666	-0.1920882562	0.9536727156
	r	-3.8872487587	10.3025699409	11.0115235230

Table 6. Numerical values of coefficients and rational terms for + + + + and + + + - helicities of $ggggh$ at kinematic point, (D.1).

In addition the values of the colour-ordered amplitudes after substitution of all the scalar integrals and including the rational terms are,

$$\begin{aligned}
 H^{1234}(1^+, 2^+, 3^+, 4^+; h) &= +29.24088185 - 46.63892079 i, \\
 |H^{1234}(1^+, 2^+, 3^+, 4^+; h)| &= 55.04741687 \\
 H^{1234}(1^+, 2^+, 3^+, 4^-; h) &= -28.10008864 + 9.836858255 i, \\
 |H^{1234}(1^+, 2^+, 3^+, 4^-; h)| &= 29.77211383 \\
 H^{1234}(1^+, 2^-, 3^+, 4^-; h) &= +4.580787288 + 7.498254006 i, \\
 |H^{1234}(1^+, 2^-, 3^+, 4^-; h)| &= 8.786775593 \\
 H^{1234}(1^+, 2^+, 3^-, 4^-; h) &= +0.369177073 - 1.815728344 i, \\
 |H^{1234}(1^+, 2^+, 3^-, 4^-; h)| &= 1.852879146
 \end{aligned} \tag{D.2}$$

Helicities	Coefficient	Real Part	Imaginary Part	Absolute Value
+ - + -	$d_{4 \times 3 \times 21}$	-7.6953556408	-6.4085129013	10.0143664825
	$d_{1 \times 23 \times 4}$	-2.3752436126	1.8890031582	3.0348171528
	$d_{1 \times 2 \times 3}$	-14.9620839628	-39.7624750054	42.4843309359
	$c_{3 \times 4}$	-0.0785670511	0.1424216217	0.1626551563
	$c_{2 \times 34}$	5.9965313107	-4.5593453199	7.5329952546
	$c_{12 \times 34}$	-22.6495761599	21.1031361652	30.9571584004
	$c_{1 \times 234}$	-9.2908649214	-2.0570320613	9.5158579166
	b_{34}	0.0682767006	0.0433975227	0.0809015007
	b_{234}	2.3825660060	-0.8884219110	2.5428162074
	b_{1234}	-4.2679248420	3.0120566624	5.2237599288
	r	2.3606586680	0.8025702116	2.4933568319
+ + - -	$d_{1 \times 2 \times 34}$	-0.0125227093	-0.5196169994	0.5197678754
	$d_{1 \times 4 \times 32}$	8.4132295790	-459.7920528912	459.8690186715
	$d_{2 \times 34 \times 1}$	62.3890431832	-51.7566409441	81.0625844095
	$d_{1 \times 23 \times 4}$	-3.4240833144	4.6410747884	5.7674883386
	$d_{1 \times 2 \times 3}$	-5.4385640586	-6.5811803202	8.5375589853
	$c_{2 \times 3}$	-41.9249189131	-23.3819669075	48.0043248295
	$c_{1 \times 23}$	-1036.7850502032	-480.4884415677	1142.7127297817
	$c_{23 \times 41}$	1080.7316959848	740.3414428401	1309.9948284984
	$c_{1 \times 234}$	22.0281339875	-305.0638529285	305.8581256899
	b_{23}	-5.3092820284	-9.1916846550	10.6148736429
	b_{234}	-0.8234906782	5.4869406052	5.5483920285
	b_{1234}	26.7533037866	15.0643874405	30.7030134100
	r	1.3340510134	0.8053633104	1.5583010518

Table 7. Numerical values of coefficients and rational terms for + - + - and + + - - helicities of $gggh$ at kinematic point, (D.1).

$$\begin{aligned}
H^{34}(1^+, 2^-, 3^+, 4^+; h) &= -8.998796972 - 13.02970981 i, \\
|H^{34}(1^+, 2^-, 3^+, 4^+; h)| &= 15.83514081 \\
H^{34}(1^+, 2^-, 3^-, 4^+; h) &= -3.850947633 + 1.791151530 i, \\
|H^{34}(1^+, 2^-, 3^-, 4^+; h)| &= 4.247119197 \\
H^{34}(1^+, 2^-, 3^+, 4^-; h) &= -0.412185752 + 7.682564596 i, \\
|H^{34}(1^+, 2^-, 3^+, 4^-; h)| &= 7.693613966
\end{aligned} \tag{D.3}$$

$$\begin{aligned}
H^{4q}(1^+, 2^-, 3^+, 4^-; h) &= 0.620045806 + 4.7030845622 i, \\
|H^{4q}(1^+, 2^-, 3^+, 4^-; h)| &= 4.743781319
\end{aligned} \tag{D.4}$$

Helicities	Coefficient	Real Part	Imaginary Part	Absolute Value
+ − + +	$d_{4 \times 3 \times 21}$	4.0685161820	−4.0500901147	5.7407363517
	$d_{4 \times 21 \times 3}$	425.5033072909	1294.6650310348	1362.7951449502
	$c_{3 \times 21}$	−73.7590711176	−242.0029027576	252.9936867102
	$c_{12 \times 34}$	14.7023801790	−6.9563781545	16.2650293561
	$c_{4 \times 123}$	32.6756691373	92.0151860555	97.6447326711
	$c_{3 \times 412}$	87.6696567417	249.0681532719	264.0471807982
	b_{12}	1.4197098901	0.3520351648	1.4627046624
	b_{124}	−0.5233075583	−1.0058683561	1.1338527023
	b_{123}	−0.7954324907	−2.6428744805	2.7599815882
	b_{1234}	−0.1009698411	3.2967076718	3.2982535351
	r	−5.4119652752	0.7153882121	5.4590428130
+ − − +	$d_{4 \times 3 \times 21}$	1.0782715488	−4.7280903169	4.8494852900
	$d_{4 \times 21 \times 3}$	13.3402061977	−3.4340877490	13.7751246842
	$c_{3 \times 21}$	−0.8475265952	2.4513967233	2.5937708504
	$c_{3 \times 4}$	0.0094395944	0.0336984573	0.0349955993
	$c_{12 \times 34}$	−3.7289304305	−10.8894201371	11.5101864919
	$c_{4 \times 123}$	1.7886984296	−1.6881229718	2.4595123988
	$c_{3 \times 412}$	1.8223566626	0.0772529014	1.8239933708
	b_{12}	−0.1514791222	−0.2049070241	0.2548191770
	b_{34}	0.0053272676	0.1024178437	0.1025562991
	b_{124}	0.2906270969	−0.1175213822	0.3134890504
	b_{123}	0.6201039140	0.1805231188	0.6458463135
	b_{1234}	−0.7645791563	0.0394874437	0.7655981613
	r	−0.2305425036	1.4755649890	1.4934663983
+ − + −	$c_{4 \times 123}$	0.0795879764	−1.9432491013	1.9448782264
	b_{123}	−1.0956687877	0.8092787161	1.3621388082

Table 8. Numerical values of the needed coefficients and rational terms for + − + +, + − − + and + − + − helicities of the $\bar{q}qggh$ process at kinematic point, (D.1).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, *Phys. Rev. Lett.* **114** (2015) 212001 [[arXiv:1503.06056](https://arxiv.org/abs/1503.06056)] [[INSPIRE](#)].
- [2] B. Mistlberger, *Higgs boson production at hadron colliders at N^3LO in QCD*, *JHEP* **05** (2018) 028 [[arXiv:1802.00833](https://arxiv.org/abs/1802.00833)] [[INSPIRE](#)].

- [3] J.M. Campbell, R.K. Ellis and G. Zanderighi, *Next-to-Leading order Higgs + 2 jet production via gluon fusion*, *JHEP* **10** (2006) 028 [[hep-ph/0608194](#)] [[INSPIRE](#)].
- [4] G. Cullen et al., *Next-to-Leading-Order QCD Corrections to Higgs Boson Production Plus Three Jets in Gluon Fusion*, *Phys. Rev. Lett.* **111** (2013) 131801 [[arXiv:1307.4737](#)] [[INSPIRE](#)].
- [5] S.P. Jones, M. Kerner and G. Luisoni, *Next-to-Leading-Order QCD Corrections to Higgs Boson Plus Jet Production with Full Top-Quark Mass Dependence*, *Phys. Rev. Lett.* **120** (2018) 162001 [[arXiv:1802.00349](#)] [[INSPIRE](#)].
- [6] CMS collaboration, *Inclusive search for a highly boosted Higgs boson decaying to a bottom quark-antiquark pair*, *Phys. Rev. Lett.* **120** (2018) 071802 [[arXiv:1709.05543](#)] [[INSPIRE](#)].
- [7] R.K. Ellis, I. Hinchliffe, M. Soldate and J.J. van der Bij, *Higgs Decay to $\tau^+\tau^-$: A Possible Signature of Intermediate Mass Higgs Bosons at the SSC*, *Nucl. Phys. B* **297** (1988) 221 [[INSPIRE](#)].
- [8] U. Baur and E.W.N. Glover, *Higgs Boson Production at Large Transverse Momentum in Hadronic Collisions*, *Nucl. Phys. B* **339** (1990) 38 [[INSPIRE](#)].
- [9] V. Del Duca, W. Kilgore, C. Oleari, C. Schmidt and D. Zeppenfeld, *Gluon fusion contributions to $H + 2$ jet production*, *Nucl. Phys. B* **616** (2001) 367 [[hep-ph/0108030](#)] [[INSPIRE](#)].
- [10] T. Neumann and C. Williams, *The Higgs boson at high p_T* , *Phys. Rev. D* **95** (2017) 014004 [[arXiv:1609.00367](#)] [[INSPIRE](#)].
- [11] G. Cullen et al., *GOSAM-2.0: a tool for automated one-loop calculations within the Standard Model and beyond*, *Eur. Phys. J. C* **74** (2014) 3001 [[arXiv:1404.7096](#)] [[INSPIRE](#)].
- [12] V. Hirschi and O. Mattelaer, *Automated event generation for loop-induced processes*, *JHEP* **10** (2015) 146 [[arXiv:1507.00020](#)] [[INSPIRE](#)].
- [13] N. Greiner, S. Höche, G. Luisoni, M. Schönherr and J.-C. Winter, *Full mass dependence in Higgs boson production in association with jets at the LHC and FCC*, *JHEP* **01** (2017) 091 [[arXiv:1608.01195](#)] [[INSPIRE](#)].
- [14] F. Buccioni et al., *OpenLoops 2*, *Eur. Phys. J. C* **79** (2019) 866 [[arXiv:1907.13071](#)] [[INSPIRE](#)].
- [15] R.K. Ellis and S. Seth, *On Higgs boson plus gluon amplitudes at one loop*, *JHEP* **11** (2018) 006 [[arXiv:1808.09292](#)] [[INSPIRE](#)].
- [16] J.M. Lindert, K. Kudashkin, K. Melnikov and C. Wever, *Higgs bosons with large transverse momentum at the LHC*, *Phys. Lett. B* **782** (2018) 210 [[arXiv:1801.08226](#)] [[INSPIRE](#)].
- [17] T. Neumann, *NLO Higgs+jet production at large transverse momenta including top quark mass effects*, *J. Phys. Comm.* **2** (2018) 095017 [[arXiv:1802.02981](#)] [[INSPIRE](#)].
- [18] R.K. Ellis and G. Zanderighi, *Scalar one-loop integrals for QCD*, *JHEP* **02** (2008) 002 [[arXiv:0712.1851](#)] [[INSPIRE](#)].
- [19] A. van Hameren, *OneLOop: For the evaluation of one-loop scalar functions*, *Comput. Phys. Commun.* **182** (2011) 2427 [[arXiv:1007.4716](#)] [[INSPIRE](#)].

- [20] S. Carrazza, R.K. Ellis and G. Zanderighi, *QCDLoop: a comprehensive framework for one-loop scalar integrals*, *Comput. Phys. Commun.* **209** (2016) 134 [[arXiv:1605.03181](#)] [[INSPIRE](#)].
- [21] D.B. Melrose, *Reduction of Feynman diagrams*, *Nuovo Cim.* **40** (1965) 181 [[INSPIRE](#)].
- [22] W.L. van Neerven and J.A.M. Vermaseren, *Large loop integrals*, *Phys. Lett.* **137B** (1984) 241 [[INSPIRE](#)].
- [23] Z. Bern, L.J. Dixon and D.A. Kosower, *Dimensionally regulated pentagon integrals*, *Nucl. Phys. B* **412** (1994) 751 [[hep-ph/9306240](#)] [[INSPIRE](#)].
- [24] Z. Bern and A.G. Morgan, *Massive loop amplitudes from unitarity*, *Nucl. Phys. B* **467** (1996) 479 [[hep-ph/9511336](#)] [[INSPIRE](#)].
- [25] L.J. Dixon, *A brief introduction to modern amplitude methods*, in *Proceedings, 2012 European School of High-Energy Physics (ESHEP 2012): La Pommeraye, Anjou, France, June 06–19, 2012*, pp. 31–67, [arXiv:1310.5353](#) [[DOI](#)] [[INSPIRE](#)].
- [26] R.K. Ellis, Z. Kunszt, K. Melnikov and G. Zanderighi, *One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts*, *Phys. Rept.* **518** (2012) 141 [[arXiv:1105.4319](#)] [[INSPIRE](#)].
- [27] R. Britto, F. Cachazo and B. Feng, *Generalized unitarity and one-loop amplitudes in $N = 4$ super-Yang-Mills*, *Nucl. Phys. B* **725** (2005) 275 [[hep-th/0412103](#)] [[INSPIRE](#)].
- [28] D. Forde, *Direct extraction of one-loop integral coefficients*, *Phys. Rev. D* **75** (2007) 125019 [[arXiv:0704.1835](#)] [[INSPIRE](#)].
- [29] P. Mastrolia, *Double-Cut of Scattering Amplitudes and Stokes' Theorem*, *Phys. Lett. B* **678** (2009) 246 [[arXiv:0905.2909](#)] [[INSPIRE](#)].
- [30] W.B. Kilgore, *One-loop Integral Coefficients from Generalized Unitarity*, [arXiv:0711.5015](#) [[INSPIRE](#)].
- [31] S. Davies, *One-Loop QCD and Higgs to Partons Processes Using Six-Dimensional Helicity and Generalized Unitarity*, *Phys. Rev. D* **84** (2011) 094016 [[arXiv:1108.0398](#)] [[INSPIRE](#)].
- [32] S.D. Badger, *Direct Extraction Of One Loop Rational Terms*, *JHEP* **01** (2009) 049 [[arXiv:0806.4600](#)] [[INSPIRE](#)].
- [33] A. Hodges, *Eliminating spurious poles from gauge-theoretic amplitudes*, *JHEP* **05** (2013) 135 [[arXiv:0905.1473](#)] [[INSPIRE](#)].
- [34] S. Badger, H. Frellesvig and Y. Zhang, *A Two-Loop Five-Gluon Helicity Amplitude in QCD*, *JHEP* **12** (2013) 045 [[arXiv:1310.1051](#)] [[INSPIRE](#)].
- [35] S. Badger, *Automating QCD amplitudes with on-shell methods*, *J. Phys. Conf. Ser.* **762** (2016) 012057 [[arXiv:1605.02172](#)] [[INSPIRE](#)].
- [36] H.B. Hartanto, S. Badger, C. Brønnum-Hansen and T. Peraro, *A numerical evaluation of planar two-loop helicity amplitudes for a W-boson plus four partons*, *JHEP* **09** (2019) 119 [[arXiv:1906.11862](#)] [[INSPIRE](#)].

- [37] G. Laurentis and D. Maître, *Extracting analytical one-loop amplitudes from numerical evaluations*, *JHEP* **07** (2019) 123 [[arXiv:1904.04067](#)] [[INSPIRE](#)].
- [38] G. Ossola, C.G. Papadopoulos and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl. Phys. B* **763** (2007) 147 [[hep-ph/0609007](#)] [[INSPIRE](#)].
- [39] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *Fusing gauge theory tree amplitudes into loop amplitudes*, *Nucl. Phys. B* **435** (1995) 59 [[hep-ph/9409265](#)] [[INSPIRE](#)].
- [40] A.G. Morgan, *Second order fermions in gauge theories*, *Phys. Lett. B* **351** (1995) 249 [[hep-ph/9502230](#)] [[INSPIRE](#)].
- [41] S. Dawson and R.P. Kauffman, *Higgs boson plus multi-jet rates at the SSC*, *Phys. Rev. Lett.* **68** (1992) 2273 [[INSPIRE](#)].
- [42] R.P. Kauffman, S.V. Desai and D. Risal, *Production of a Higgs boson plus two jets in hadronic collisions*, *Phys. Rev. D* **55** (1997) 4005 [Erratum *ibid.* **D 58** (1998) 119901] [[hep-ph/9610541](#)] [[INSPIRE](#)].
- [43] S. Badger, E.W. Nigel Glover, P. Mastrolia and C. Williams, *One-loop Higgs plus four gluon amplitudes: Full analytic results*, *JHEP* **01** (2010) 036 [[arXiv:0909.4475](#)] [[INSPIRE](#)].
- [44] R.K. Ellis, W.T. Giele, Z. Kunszt and K. Melnikov, *Masses, fermions and generalized D-dimensional unitarity*, *Nucl. Phys. B* **822** (2009) 270 [[arXiv:0806.3467](#)] [[INSPIRE](#)].
- [45] J.M. Campbell, R.K. Ellis and C. Williams, *Vector boson pair production at the LHC*, *JHEP* **07** (2011) 018 [[arXiv:1105.0020](#)] [[INSPIRE](#)].
- [46] J.M. Campbell, R.K. Ellis and W.T. Giele, *A Multi-Threaded Version of MCFM*, *Eur. Phys. J. C* **75** (2015) 246 [[arXiv:1503.06182](#)] [[INSPIRE](#)].
- [47] J. Campbell and T. Neumann, *Precision Phenomenology with MCFM*, *JHEP* **12** (2019) 034 [[arXiv:1909.09117](#)] [[INSPIRE](#)].
- [48] S.D. Badger, E.W.N. Glover, V.V. Khoze and P. Švrček, *Recursion relations for gauge theory amplitudes with massive particles*, *JHEP* **07** (2005) 025 [[hep-th/0504159](#)] [[INSPIRE](#)].
- [49] R. Britto and B. Feng, *Unitarity cuts with massive propagators and algebraic expressions for coefficients*, *Phys. Rev. D* **75** (2007) 105006 [[hep-ph/0612089](#)] [[INSPIRE](#)].